

## Durham Research Online

---

### Deposited in DRO:

04 August 2016

### Version of attached file:

Accepted Version

### Peer-review status of attached file:

Peer-reviewed

### Citation for published item:

Hillier, J.K. and Kougiumtzoglou, I.A. and Stokes, C.R. and Smith, M.J. and Clark, C. and Spagnolo, M.S. (2016) 'Exploring explanations of subglacial bedform sizes using statistical models.', PLoS ONE., 11 (7). e0159489.

### Further information on publisher's website:

<http://dx.doi.org/10.1371/journal.pone.0159489>

### Publisher's copyright statement:

Copyright: © 2016 Hillier et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Additional information:

## Use policy

---

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

# Exploring Explanations of Subglacial Bedform Sizes Using Statistical Models

Hillier<sup>1\*</sup>, J.K., Kogioumtzoglou<sup>2</sup>, I. A., Stokes<sup>3</sup>, C. R., Smith<sup>4</sup>, M. J., Clark<sup>5</sup>, C. D., Spagnolo<sup>6</sup> M. S.

<sup>1</sup>Department of Geography, Loughborough University, UK.

<sup>2</sup>Department of Civil Engineering and Engineering Mechanics, Columbia University, USA.

<sup>3</sup>Department of Geography, Durham University, UK.

<sup>4</sup>School of Geography, Earth and Environment, Kingston University, UK.

<sup>5</sup>Department of Geography, University of Sheffield, UK.

<sup>6</sup>School of Geosciences, University of Aberdeen, UK.

\*Corresponding Author

E-mail: J.Hillier@lboro.ac.uk (JH)

## Abstract

Sediments beneath modern ice sheets exert a key control on their flow, but are largely inaccessible except through geophysics or boreholes. In contrast, palaeo-ice sheet beds are accessible, and typically characterised by numerous bedforms. However, the interaction between bedforms and ice flow is poorly constrained and it is not clear how bedform sizes might reflect ice flow conditions. To better understand this link we present a first exploration of a variety of statistical models to explain the size distribution of some common subglacial bedforms (i.e., drumlins, ribbed moraine, MSGSL). By considering a range of models, constructed to reflect key aspects of the physical processes, it is possible to infer that the size distributions are most effectively explained when the dynamics of ice-water-sediment

interaction associated with bedform growth is fundamentally random. A ‘*stochastic instability*’ (SI) model, which integrates random bedform growth and shrinking through time with exponential growth, is preferred and is consistent with other observations of palaeo-bedforms and geophysical surveys of active ice sheets. Furthermore, we give a proof-of-concept demonstration that our statistical approach can bridge the gap between geomorphological observations and physical models, directly linking measurable size-frequency parameters to properties of ice sheet flow (e.g., ice velocity). Moreover, statistically developing existing models as proposed allows quantitative predictions to be made about sizes, making the models testable; a first illustration of this is given for a hypothesised repeat geophysical survey of bedforms under active ice. Thus, we further demonstrate the potential of size-frequency distributions of subglacial bedforms to assist the elucidation of subglacial processes and better constrain ice sheet models.

## 1. Introduction

Observations of palaeo-ice sheet beds show sediment that is commonly organized into subglacial bedforms (e.g., drumlins), whose shape or occurrence is thought to reflect ice flow conditions [1–3]. Concurrently, these bedforms are also thought to modulate ice flow characteristics, such as velocity ( $v$ ) through their effect on subglacial hydrology, basal friction and roughness [4–7]. In short, there is likely an association between bedform morphology and the behaviour of the ice-sediment-water system that drives their formation.


Recently, geophysical observations from an Antarctic ice stream have revealed bed conditions [8–10] and bedforms that evolve, grow, and shrink on sub-decadal timescales [11–14]. However, these observations are logistically challenging and so limited to relatively few bedforms at one site [13,14]. In contrast, palaeo-bedforms are abundant (i.e.,  $> 100,000$ s) and widespread, but it is more challenging to link them securely to processes at the ice sheet bed. Thus, our understanding of the processes occurring beneath contemporary ice sheets is

53 incomplete, with some fundamental questions largely unanswered, e.g., how do bedforms  
54 grow, evolve their shape (e.g., elongate), regulate sediment flux, and interact with basal  
55 conditions such as 'sticky spots' [e.g., 15]?


56 Size-frequency statistics of observed groups of bedforms thought to be genetically linked  
57 (Fig. 1), known as 'flow sets' [e.g., 16] or 'fans' [17], may provide an additional powerful  
58 constraint on such questions [e.g., 18,19]. However, these statistics are under-exploited, and  
59 factors such as the shape of the frequency distribution have been given only limited attention.  
60 Distribution shape has been neglected as a constraint because the current conceptual and  
61 physics-based models do not predict bedform size-frequency distributions. The potential to  
62 act as a constraint arises because not all conceptual or physics-based models [e.g., 20,21]  
63 explaining bedform growth will replicate the observed sizes. Statistical models [19,22],  
64 however, have the potential to predict bedform sizes as a combined product of key aspects of  
65 the physical process: antecedent bedform-scale topography, growth rate (e.g., exponential),  
66 and the timing of growth. Fig. 2 illustrates size distributions produced by a variety of  
67 statistical models, some of which are consistent with the shape of observed distributions and  
68 some are not.

69 Hillier et al. [19] first proposed a conceptual model to explain subglacial bedforms' size-  
70 distributions, in which ice-sediment-water interaction creating bedforms is fundamentally  
71 stochastic. Specifically, to explain an exponential tail to the size-distribution, this model  
72 suggests that bedform growth processes may be a convolution of randomness with simple  
73 rules about their rate of growth; analogous models of 'self-organized criticality' are used to  
74 explain power-law distributions [23,24]. The subglacial model draws upon ideas of  
75 probabilistic sediment transport [i.e., 25] and an analogy to fluvial bedforms whose heavy-  
76 tailed size-distributions are thought to originate through growth in the presence of random  
77 fluctuations associated with turbulent flow [26–30]. As a concept this is consistent with the  
78 geophysical observations in Antarctica, but does not necessarily exclude either ice-till [e.g.,  
79 20] or meltwater [e.g., 21] bedform growth models. Fowler et al. [22] formalized a first

statistical model of bedform sizes, investigating explanations for the particular case of a log-normal approximation to the observed size-distribution under the assumption of exponential growth without shrinking. This paper, to better understand how bedform sizes might reflect ice flow conditions, re-formulates and develops Fowler's statistical model and creates a new range of other models. This variety of models is a first exploration of the possibilities and allows, by putting each model in context, an assessment of its relative plausibility.



**Fig. 1 Size-frequency data and statistical distributions fitted to them.** a) to c) Normalised histograms of observed drumlin attributes on semi-log plots (black dots), to which selected statistical distributions are fitted and plotted as probability density functions (pdfs); exponential distribution (solid blue line); gamma distribution (dashed line) ( $\alpha_{\text{obs}}$ ,  $\beta_{\text{obs}}$ ) [19]; log-normal (dotted line) ( $\mu_{\text{obs}}$ ,  $\sigma_{\text{obs}}$ ) [22]. Fits to obtain the distribution parameters, shown as Greek letters, are performed using estimators (e.g., maximum likelihood) as detailed in Appendix B. Data source and number of observed bedforms  $n$  are indicated on the plots; country-wide UK data (Fig. 8 in [16] and Fig. 5 in [31]) (black) and a well-studied sub-set (grey) of this [32] are used. d) The typical shape; there are few small bedforms, a modal peak above this forming a 'roll-over', and an approximately exponential tail of frequencies decreasing towards the largest sizes.



**Fig. 2 Illustrative size-frequency distributions from statistical growth models.** Semi-log frequency plot illustrating a variety of size-frequency distributions of bedforms predicted by different types of statistical growth model. They are each governed by arguably plausible glaciological or statistical assumptions (see text for models): Dirac delta function (dot-dash line is Model 1, denoted M1); uniform distribution (dotted line e.g., M4); exponential (solid line e.g., M8); log-normal (dashed line e.g., M7). The power of this size-frequency data as a constraint is that only a sub-set of models produces distributions reasonably approximating observed data (e.g., Fig. 1).

The paper begins by describing the size-frequency observations of bedforms (i.e., drumlins, ribbed moraine, MSGSL), then outlines the terminology and defines a conceptual framework necessary for statistically modelling the evolution of sets of such subglacial bedforms. It then builds new statistical models, which are evaluated and discussed in light of observational evidence, internal consistency, and their implications for theories of bedform growth and the ice-water-sediment system under ice sheets. In addition, the models are shown to make distinctive predictions that could be tested should a geophysical survey under active ice [i.e., 13] be repeated. Because growth in bedform height ( $H$ ) underlies most physical modelling [e.g., 20,33,34] the models are initially developed for height, but with implications for width ( $W$ ) and length ( $L$ ) also discussed.

## 2. Size Observations

Fig. 1 illustrates typical size-frequency statistics of observed groups of subglacial bedforms. Distribution shapes are similar across bedform types (i.e., drumlins, MSGSL, ribbed

moraine), mappers and regions (e.g. UK, Canada, Sweden) [19]. Although a selection of statistical distributions could be fitted to bedform size data [e.g., 26], subglacial bedform sizes have been found to be reasonably approximated as having a log-normal shape [22,35,36] or as being exponential above their mode [19]. Large compilations of bedforms ( $n > 10,000$ ) [e.g., 16] more precisely constrain their size distribution than smaller ones as uncertainty in sampling is reduced, but almost certainly represent the aggregation of a range of subglacial conditions. As such, the size distributions of large compilations may simply represent the statistical effects of aggregating samples rather than anything to do with ice flow. It is therefore important to note that the same distribution shape and spread of sizes is still apparent within flow-sets comprising 100-200 bedforms (Fig. 1, grey lines) that likely represent something about glaciological conditions at a particular location in space and time.

The parameters listed in Fig. 1 for the best-fitting gamma ( $\alpha$ ,  $\beta$ ) and log-normal ( $\mu$ ,  $\sigma$ ) distributions are obtained by method of moment and maximum likelihood estimators as described in Appendix B. Country-wide UK data in Fig. 1 are, quite deliberately, values digitised from plots in the original papers [16,31]. This is done to demonstrate that the published archive of size-distributions can be usefully re-assessed in light of statistical models. Parameters calculated from digitized values typically differ little from those used to construct the original plots (e.g.,  $<3\%$  for  $\mu$  and  $\sigma$ ). Furthermore, the data of Hillier and Smith [32] show that parameter values are similar when calculated from either counts within size bins or from the individual underlying data (e.g., variations  $<7\%$  for  $\mu$  and  $\sigma$ ). Importantly, patterns in relative values (e.g.,  $\sigma_H > \sigma_W > \sigma_L$ ) are robustly unchanged for all parameters, and the differences between their values (e.g., for  $H$  vs.  $W$ ) are always substantially larger than uncertainties caused by the method used to derive the parameter values (see Supporting Information).

Initially, the parameters are simply empirical descriptors of the shape of the size-frequency distributions; it is statistical models of bedform growth that potentially allow the parameters to be considered in terms of subglacial processes. A conceptual framework is now created,

which outlines the elements necessary to formulate statistical models that might explain the observed size-frequency distributions.

### 3. Conceptual Framework

Firm and direct observational constraints on how glacial bedforms are formed have proved challenging to obtain. However, to formalise statistical models, a framework is needed. Geophysical surveys [11,13], sediment flux estimates [37], and geometric arguments [38] indicate that forms entirely composed of sediment could arise over ~10s-100s years, and certainly within one ice flow event [e.g., 39,40]. Thus, modelling can start by considering one flow episode. However, substantial elements of the processes at work remain unclear. How do bedforms initiate? Do initial sizes determine final ones? Is growth exponential with time, characteristic of linear instability? Is growth continuous or discrete, and monotonic or fluctuating, over time? Are bedforms in equilibrium with ice flow? It is not practical to model all views held on these questions, so these topics are introduced in order to highlight the choices made in constructing the statistical models.


#### 3.1. Bedform initiation: growth and location

Entirely bedrock bedforms exist, and require an erosional mechanism [e.g., 41]. The majority, however, appear to be composed mainly or entirely of glacially-derived sediment (i.e., till) [42,43] requiring a mechanism for an origin from a till sheet [e.g., see 44]; this could involve erosion, deposition or redistribution or a combination of any of these processes [e.g., 45]. Subglacial bedforms might decrease in height from some set of progenitor forms [e.g., 46]. Alternatively, if sculpted from a relatively flat surface, they must (as a net effect over a period of time) increase their amplitude or ‘grow’ [e.g., 20]. This paper considers a sub-set of statistical models of bedform genesis in which bedforms undergo net growth, including models that incorporate periods where bedforms are stable or shrink. The mechanism of net growth may be till deformation [e.g., 47,48] but, especially in light of studies into the



175 size distribution of fluvial scours [e.g., 49], the statistical models may also apply to  
 176 conceptual models of the ice-sediment-water system governed by erosion or scour by  
 177 meltwater [e.g., 21,50,51].

178 It is known that bedforms occur more densely in some places than others, creating  
 179 patchiness on a scale of 10-100s of km [e.g., 52,53]. ‘Patches’ defined in this way encompass  
 180 numerous individual bedforms, which are typically 0.1-10 km in horizontal extent. Thus,  
 181 meso-scale ( $\sim 10$ -100s km) ‘patches’ are envisaged for the statistical models (Fig. 3), which  
 182 contain a statistically useful number (i.e., 1,2,3 ....  $j$ ) of bedforms linked to relatively local  
 183 conditions (black dots) that grow in height (i.e.,  $H$ ). The premise of using patches as defined  
 184 is consistent with the idea of spatio-temporally variable mosaics of stable and deforming bed  
 185 conditions; this is based on observations of exposed till [54,55], but also consistent with  
 186 geophysical studies that have revealed variable bed conditions [9,10]. Spatial variation in  
 187 conditions is also postulated in bedform models that invoke meltwater [56].



188 **Fig. 3 Conceptualisation of how flow-sets of bedforms grow.** a) Cross-hatched area is a  
 189 meso-scale flow-set ( $\sim 10$ -100 km) or ‘patch’ of deformable or erodible subglacial material  
 190 subjected to conditions conducive to a flow set of bedforms arising in locations illustrated by  
 191 black dots. Within this, bedforms from 1 to  $j$ , where  $j$  is any integer, change in amplitude  
 192 through erosion, deposition, or redistribution. b) A potential, illustrative, sequence of growth  
 193 for one bedform (number  $j$ ) through time (dashed line), accompanied by selected silhouettes  
 194 representing vertical cross-sections; a shrinking rate of zero (i.e., stasis) is valid within the  
 195 illustration.  
 196

### 3.2. Growth style: deterministic versus probabilistic

‘*Deterministic*’ growth is where proto-bedforms of a given size and shape always evolve similarly with time to a predictable final morphology; i.e., initial conditions lead uniquely to a final configuration. ‘*Probabilistic*’ growth is where random variability through time (i.e., dynamics) causes individual bedforms to evolve unpredictably or ‘*stochastically*’, but combine to produce predictable flow set statistics [e.g., 18,57]. In the non-turbulent conditions of ice flow, such variability is likely to arise from time-varying boundary conditions in the coupled ice-sediment-water system (e.g., water incursions, floods, basal stick-slip events) [58–61] or interactions between bedforms [62] perhaps by ice rheology inducing lateral stresses [e.g., 63,64]. Combining this with the observed range of time-scales on which ice flow fluctuates (i.e., days to decades) [e.g., 60,65–74], and by analogy with established ideas in fluvial and aeolian environments [e.g., 25,28–30,57,62], gives a picture of potentially pervasive randomness through time in subglacial sediment transport (i.e., flux)[19]. Either deterministic or probabilistic growth can be readily incorporated into statistical models.

### 3.3. Growth rate

Bedform growth predicted by physics-based models proceeds at a rate that has an expected characteristic mathematical form. If models relate till flux to the thickness of the till body and an unconnected ‘field’ variable, such as basal shear stress ( $\tau$ ), that can vary in space [e.g., 20,75,76], growth of  $H$  is initially linear with time at a constant rate ( $k$ ). In this regard  $H$  is governed by the ordinary differential equation (ODE)

$$\frac{dH}{dt} = k \quad \text{Eq. 1}$$

in conjunction with the initial condition

$$H(t_i) = H_i. \quad \text{Eq. 2}$$

Integrating Eq. 1 analytically, considering the initial condition, and for final height denoting  $H(t_f) = H_f$ , yields Eq. 3.

$$H_f = H_i + k(t_f - t_i) \quad \text{Eq. 3}$$

If, on the other hand, models contain positive linear feedback between bedform and ‘field’ (Eq. 4), this results in a physical instability in the sediment-ice system and growth is initially exponential with time (Eq. 5) [e.g., 20,33]. Thus, the term ‘instability’ has been adopted to describe this class of sediment growth model. Note that the term instability is used in this way in this paper and not as strictly defined in the mathematical field of stability theory related to dynamics.

In this regard, where physical processes are thought to be approximated by linear feedback,  $H$  is governed by the ODE

$$\frac{dH}{dt} = kH \quad \text{Eq. 4}$$

in conjunction with the initial condition of Eq. 2. Similarly, as with Eq. 1, integrating analytically yields

$$H_f = H_i e^{k(t_f - t_i)} \quad \text{Eq. 5}$$

It is entirely plausible that growth does not continue according to either of these simple rate laws, perhaps because of ‘shock formation’ as  $H$  increases, which is when a subglacial bedform is dramatically altered after an ice-free cavity is generated on its lee side [e.g., 77,78]. The statistical models proposed below focus on the simple rate laws as it is not yet even well determined which of these might apply [cf. 79,80,81]. The models are, however, presented initially in terms of time spent growing so that they can be readily adapted for other rate laws if required in the future.

### 3.4. Continuous process versus discrete events

If bedform growth is viewed as a continuous property extending over a finite time period [e.g., 20,48,79] then at any time, and for finite proportions of it, bedforms either grow or shrink. In contrast, and by analogy with other environments [e.g., 82,83], the creation of each bedform may occur through discrete sediment flux '*events*', each of which might affect several proximal bedforms. However, if events affect only sub-areas of a patch and are randomly located, their impacts upon each bedform will appear as a series of independent trials through time [22], analogous to continuous variability. Thus, and particularly because analogies between the continuous and discrete mathematics exist [e.g., 84], either a continuous or discrete modelling approach remains valid.

### 3.5. Transient versus equilibrium growth


The length of time over which a flow-set develops is not well constrained. It is therefore necessary to introduce into this framework the concept of '*transient*' flow-set growth within a time window, between an initial time ( $t_i$ ) and a final time ( $t_f$ ). Pre-equilibrium or transient growth is where the statistics of a flow-set evolve over time, continue to evolve, and would have continued to evolve further if the conditions for growth had persisted. This contrasts to stable long-term '*equilibrium*' behaviour in which the statistical characteristics of a flow set stabilise. Equilibrium is actively sought in fluvial experimentation [e.g., 26] and has been implicitly invoked to infer ice properties; for example, assumed equilibrium is implicit when arguing that bedform elongation is related to ice velocity, rather than duration of flow [e.g., 3,85]. Bedforms that develop slowly with respect to changes in ice flow conditions at the flow-set scale (~10-100 km) will have pre-equilibrium transient statistics, whilst forms evolving much more rapidly than patch-scale flow changes could attain equilibrium. Which behaviour predominates amongst glacial bedforms is not yet known. Thus, statistical models containing both behaviours are permitted and explored here.

## 4. Methods

274 To better understand how bedform sizes might reflect formative flow conditions a new  
275 range of statistical models are developed, including one that extends the model of Fowler et al  
276 [22]. This variety of models allows, by putting each model in context, an assessment of its  
277 relative plausibility. The initial mode of discrimination is by the shape of the size-frequency  
278 distribution that each model creates (e.g., Fig. 2) as compared to observations. Specifically, as  
279 also demonstrated in Fig. 1, the data are reasonably approximated by log-normal[22,35,36]  
280 and gamma distributions, and by an exponential tail above the mode [19]. Models are  
281 therefore required to generate at least one of these to be considered as potentially plausible.  
282 Models are developed analytically so that the form of the size-frequency distributions they  
283 can produce is known explicitly.

## 284 5. Models

285 The models developed here contain a number (i.e., 1,2,3 ....  $j$ ) of non-overlapping bedforms  
286 (Fig. 4a, black dots) characterised as growing independently for a time period between  $t_i$  and  
287  $t_f$  within 'meso-scale' ( $\sim 10$ -100s km) '*patches*' when an appropriate flow regime prevails.  
288 Statistical independence between bedforms is assumed as in previous statistical modelling  
289 [i.e., 22], where it is justified by randomness in the perturbing field (e.g., water influx) (see  
290 Section 3.4), although it may also be augmented by spatial randomness in rheological  
291 properties (e.g., viscosity). This is consistent with stochastic sediment flux in aeolian cellular-  
292 automata models that has yielded randomly sized, yet spatially patterned, barchan dunes  
293 [62,86]. Effective independence is also supported by analogy to extensive work in the fluvial  
294 environment where the growth of spatially ordered and self-organized bedforms is statistically  
295 described and modelled as stochastic and random [26,28,30,57,87]. We acknowledge that,  
296 with limited observational evidence, this set-up may not ultimately turn out to be correct, but  
297 it forms a useful basis to start an exploration with statistical models. Physically, activity  
298 within the patches is conceptualised as being based on multiple, rapid (i.e., sub-decadal) and  
299 random fluctuations in basal conditions that generate flow sets of bedforms.



**Fig. 4 Framework for the statistical models.** Cross-hatched area in a) is a meso-scale (~10s-100s km) 'patch' of deformable or erodible subglacial material subject to conditions conducive to a flow set of bedforms arising. b) and c) are barcode style strips for the waiting time (WT) [M10] and stochastic instability (SI) [M7] models. The strips represent the size evolution through time for one of the bedforms  $j$  in a). Specifically, the bands represent alternating 'local' (~0.1-1 km) conditions affecting  $H$ ; grey is growth, and white is shrinking or inactivity.  $k$  and  $kH$  indicate growth rate (i.e., Eq. 1 and Eq. 4). Rapid fluctuations in c) are omitted for visual clarity, analogous to a time-series recorded at low temporal resolution.

Models are numbered, so that Model 4 is denoted [M4], for example. Each includes four elements, a growth rate 'law' based upon suggestions from physical models[20,33,75,76], rules about what initial sizes are and when growth begins, and a growth style that is deterministic or uses temporal randomness. Each aspect affects the output size distribution, and the characteristics of all models are summarised in Table 1. The simplest new models created, both mathematically and conceptually, are those that do not involve stochasticity in growth through time [M1-5]. Some of these (see Table 1) can replicate size-frequency observations (Fig. 1), but require substantial *ad hoc* assumptions to do so; for instance, in M3 a log-normal antecedent size distribution is needed to create a log-normal distribution of observed sizes [i.e., M3a]. So this preliminary exploration is detailed in Appendix A, with statistical models incorporating probabilistic growth [M6-11] focussed on below.

**Table 1: Attributes of the models.** Grey shading indicates the variable changed in each group of models. See Section 3 for a discussion of the conceptual framework, which outlines the different parts that comprise the models. SI and WT in column 1 refer to the 'Stochastic

Instability' and 'Waiting Time' models, respectively. Models 1-5 are in Appendix A. The distribution shapes each model can produce are described in sections where they are developed, and acceptable approximations to observations are log-normal, gamma or exponential above the mode.

| #        | Growth Rate 'law' |      |     | Initial sizes      |         |            | Growth Style |          |         | Growth initiation timing |                                                                                       |          |       | Can explain size-frequency observations ? |
|----------|-------------------|------|-----|--------------------|---------|------------|--------------|----------|---------|--------------------------|---------------------------------------------------------------------------------------|----------|-------|-------------------------------------------|
|          | Linear            | Exp. | Any | Dirac (i.e., same) | Uniform | Log-normal | Det.         | Brownian | Poisson | Dirac (i.e., same)       | Uniform                                                                               | Gaussian | Other |                                           |
| M1       |                   |      | ✓   | ✓                  |         |            | ✓            |          |         | ✓                        |                                                                                       |          |       | X                                         |
| M2       | ✓                 |      |     |                    | ✓       |            | ✓            |          |         | ✓                        |                                                                                       |          |       | X                                         |
| M3       |                   | ✓    |     |                    | ✓       |            | ✓            |          |         | ✓                        |                                                                                       |          |       | X                                         |
| M3a      |                   | ✓    |     |                    |         | ✓          | ✓            |          |         | ✓                        |                                                                                       |          |       | ✓                                         |
| M4       | ✓                 |      |     | ✓                  |         |            | ✓            |          |         |                          | ✓                                                                                     |          |       | X                                         |
| M4a      | ✓                 |      |     | ✓                  |         |            | ✓            |          |         |                          |                                                                                       |          | ✓     | ✓                                         |
| M5       |                   | ✓    |     | ✓                  |         |            | ✓            |          |         |                          | ✓                                                                                     |          |       | X                                         |
| M5a      |                   | ✓    |     | ✓                  |         |            | ✓            |          |         |                          |                                                                                       | ✓        |       | ✓                                         |
| M6       | ✓                 |      |     | ✓                  |         |            |              | ✓        |         | ✓                        |                                                                                       |          |       | X                                         |
| M7 (SI)  |                   | ✓    |     | ✓                  |         |            |              | ✓        |         | ✓                        | i.e., for M6-11 conditions for growth of the flow-set start at a single point in time |          |       | ✓                                         |
| M8       | ✓                 |      |     | ✓                  |         |            |              |          | ✓       | ✓                        |                                                                                       |          |       | Tail; not roll-over                       |
| M9       |                   | ✓    |     | ✓                  |         |            |              |          | ✓       | ✓                        |                                                                                       |          |       | X                                         |
| M10 (WT) | ✓                 |      |     | ✓                  |         |            |              |          | ✓       | ✓                        |                                                                                       |          |       | ✓                                         |
| M11      |                   | ✓    |     | ✓                  |         |            |              |          | ✓       | ✓                        |                                                                                       |          |       | X                                         |

If ice-sediment-water interaction leading to bedform growth is fundamentally stochastic, as proposed by the conceptual model of Hillier et al. [19], then stochastic mathematical models [e.g., 88,89] may be constructed to formalise variants on this idea. Of possible types of time-series (i.e., temporal) randomness [e.g., 90], the two most standard and well-established descriptions [e.g., 91] are selected to create simple stochastic models. Models are therefore created based on 'white noise' (Brownian motion) [M6 and M7], developing that of Fowler et al. [22], and Poisson randomness [M8 to M11] as seen in natural processes such as storms impacting land [92]. Particular attention was paid to variants capable of generating distributions that have previously been fitted as approximations to the size-frequency observations (i.e., exponential, gamma, log-normal [e.g., 19,22]).

The models employ statistical derivations from texts such as Soong [93], but also use elements from stochastic processes and stochastic differential equations [e.g., 88,94]. All analytical solutions have been validated with pertinent Monte Carlo simulations utilizing 10,000 samples compatible with the statistics of the random quantities [e.g., 95].


### 5.1. Brownian motion randomness [M6 and M7]

Models M6 and M7 incorporate probabilistic growth governed by randomness of a type known by a number of names including ‘Brownian motion’, ‘white noise’, or a ‘1D random walk’ [e.g., 94]. This latter can be pictured as a drunkard in a long, thin alleyway, who either stumbles ‘forward’ or ‘back’ randomly, leading to a distribution of positions that expands with time. If each drunken step takes 1 unit of time, then the net time travelling forward will evolve exactly as distance does, starting to differ increasingly with time, spreading out or dispersing when plotted with predictable statistics: namely, a mean of  $\mu$  and standard deviation of  $\sigma$  (Fig. 5a). Analogously, if changes to a bedform continuously fluctuate between two states (i.e., growth, g, or shrinking, s) in a manner analogous to a random walk (Fig. 4c) then net time spent growing (i.e.,  $t_N(t) = \sum t_g - \sum t_s$ ) is a random variable with a ‘diffusive’ part caused by random motions that is a Gaussian or ‘normal’ distribution [94,96]. Specifically, as the size of steps tend to zero, this is described by a Wiener process denoted  $W(t)$  [88,94,97,98] and the Gaussian distribution has mean ( $\mu$ ) of 0 and variance ( $\sigma^2$ ) of  $t$  i.e.,  $\sim N(0, t)$ . Namely,  $E[W(t)] = 0$  and  $E[W^2(t)] = t$  with the property  $W(t) - W(s) \sim N(0, t - s)$  for  $t > s \geq 0$ . Statistical ‘drift’ ( $\xi$ ) where the mean of the distribution increases or decreases with time ( $\mu = \xi t$ ) can also be accounted for [e.g., 98, p462]; this can be driven by growth being more probable, namely the probability of growing ( $p$ ) being greater than 0.5. This would represent a drunkard capable of some ability to travel forward. Thus, the distribution of  $t_N(t)$  is given by Eq. 6 and illustrated in Fig. 5a as a hump that both moves or ‘drifts’ and spreads out or ‘diffuses’.

$$t_N(t) = \text{'drift'} + \text{'diffusion'} = \xi t + W(t) \quad \text{Eq. 6}$$

369





**Fig. 5 Visualisation of the relationship between a random walk, a Wiener process, and the evolving log-normal size-frequency distribution expected of bedforms in the SI model [M7].** a) Probabilities for the number of discrete steps taken in a random walk (grey circles) are distributed binomially. From Wiener's work whatever small step length is chosen these are well approximated by normal distribution (black line) of  $\mu = 0$  and  $\sigma^2 = t$  i.e., net time spent growing is a normally distributed random variable. If  $H \propto \exp(t_N)$  this defines a log-normal distribution for  $H$ . b) Height distributions evolving through the SI model [M7] as time increases for some illustrative constants.

Alternatively, the distribution of  $t_N$  created by a Wiener process with drift can be described by a stochastic differential equation (SDE) [e.g., 88,99] (Eq. 7), which integrates to Eq. 6 under the initial condition that growth starts at  $t_i$ , namely  $t_N(t_i) = 0$ ; note that this simple case can be integrated directly since the integral of  $dW(t)$  is  $W(t)$  by definition, and it is not necessary to use Itô's formula. The pdf obtained by either means is more fully expressed by writing out the equation of a Gaussian (Eq. 8) with appropriate values of the mean ( $\mu$ ) and variance ( $\sigma^2$ ) given by Eq. 9 and Eq. 10.

$$dt_N(t) = \xi dt + dW(t) \quad \text{Eq. 7}$$

$$f(t_N) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left[-\frac{1}{2} \frac{(t_N - \mu)^2}{\sigma^2}\right], -(t_f - t_i) \leq t_N \leq (t_f - t_i) \quad \text{Eq. 8}$$

$$\mu = \xi(t_f - t_i)$$

$$\sigma^2 = (t_f - t_i) \quad \text{Eq. 9}$$

$$\sigma^2 = 4[p(1-p)](t_f - t_i) \quad \text{Eq. 10}$$

Statistical drift ( $\xi$ ) caused by varying  $p$  is given by  $\xi = 2p - 1$ . This affects the mean of  $t_N$ , giving an expression for  $\mu$  as in Eq. 11. Two special cases illustrate this behaviour. Without any directional bias, namely if probability of growing and shrinking are equal with  $p = 0.5$ ,  $\xi = 0$  and no drift occurs. If all steps are in one direction, namely  $p = 0$  or  $1$ , then there is no randomness and  $\xi = \pm 1$  as is appropriate to set growth or shrinkage to a single deterministic rate. However, in the limiting case of  $\xi = \pm 1$  the distribution of  $t_N$  cannot diffuse and spread into a Gaussian, and so the spread (i.e., variance) of  $t_N$  is also demonstrably affected by  $p$ , especially near its limits of  $0$  and  $1$ . This effect is described through well-established results; the discrete Binomial distribution ( $n, p$ ) is approximated as a Normal distribution ( $\mu, \sigma^2$ ), where  $\sigma^2 = np(1-p)$  as  $n \rightarrow \infty$  [e.g., 84] (e.g., Fig. 5a). Thus, the variance of  $t_N$  in Eq. 8 is given by Eq. 12, where the factor of  $4$  arises because the step size is doubled, namely  $(-1, +1)$  in time versus  $(0, +1)$  for the Binomial, which is squared in its impact upon the variance of a random variable [e.g., 93, p81].

$$\mu = (2p - 1)(t_f - t_i) \quad \text{Eq. 11}$$

$$\sigma^2 = 4[p(1-p)](t_f - t_i) \quad \text{Eq. 12}$$

Now, it is possible to convert back from time to height, choosing whatever growth law is desired. Firstly, recognising that  $(t_f - t_i)$  in Eq. 3 and Eq. 5 is simply a specific case of net time spent growing (i.e.,  $t_N = \sum t_g - \sum t_s$ ), equations for linear and exponential growth can be re-written as in Eq. 13 and Eq. 14, respectively. Then,  $t_N$  generated by Brownian motion

randomness from Eq. 8 can be applied to the different growth rates by transformations of the random variables [e.g., Ch 5 of 93] as in the simpler models in Appendix A (e.g., using Eq. 29).

$$H_f = H_i + kt_N \quad \text{Eq. 13}$$

$$H_f = H_i e^{kt_N} \quad \text{Eq. 14}$$

First, consider growth that is linear with time (Eq. 13). This is denoted as model M6. The overall amount of time spent growing ( $t_N$ ) is normally distributed. Since  $H_f$  is a simple multiple of this, it will also be normally distributed. As above, analytically determining the pdf of  $H_f$  given the pdf of  $t_N$  is a relatively straightforward task using the standard transformation relationship. This yields Eq. 15 to Eq. 17, which describe  $H_f$  as a Gaussian drifting and diffusing as time passes; i.e., not gamma, exponential or log-normal.

$$f_{H_f}(h_f) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left[-\frac{1}{2} \frac{(h_f - \mu)^2}{\sigma^2}\right], H_i - k(t_f - t_i) \leq h_f \leq H_i + k(t_f - t_i) \quad \text{Eq. 15}$$

$$\mu = H_i + k(2p - 1)(t_f - t_i) \quad \text{Eq. 16}$$

$$\sigma^2 = k^2 4[p(1 - p)](t_f - t_i) \quad \text{Eq. 17}$$

In contrast, model M7 is formulated for growth that is exponential (Eq. 14). Since  $t_N$  is normally distributed,  $H_f$  will be log-normally distributed by definition (see Appendix A.3 ‘Variable initiation times’). This is to say that where future increase in a variable is linearly dependent on past progress (i.e., instability, Eq. 4 or Eq. 14) a log-normal distribution is produced [e.g., 25] (Eq. 18 to Eq. 20). This assertion can be verified by analytically determining the pdf of  $H_f$  in Eq. 14 given the pdf of  $t_N$  and by using the transformation relationship for random variables. Alternatively, the same result can be reached using Stochastic Differential Equations (SDEs). Indeed the form of the result using SDEs is very well established and is known as the solution of ‘*Geometric Brownian Motion*’, which is used

for purposes such as predicting stock prices [e.g., 98,100,101]. It is important to note for comparisons, however, that common treatments using SDEs do not allow  $p$  to vary from 0.5 and, instead of  $k$ , usually use as their growth constant the effective stochastic equivalent growth rate which for  $p = 0.5$  is  $\bar{k} = \xi + k^2/2$  [e.g., 101, p546].

$$f_{H_f}(h_f) = \frac{1}{\sigma h_f \sqrt{2\pi}} \exp \left[ -\frac{1}{2} \frac{(\ln(h_f) - \mu)^2}{\sigma^2} \right], H_i e^{-k(t_f - t_i)} \leq h_f \leq H_i e^{k(t_f - t_i)} \quad \text{Eq. 18}$$

$$\mu = \ln(H_i) + k(2p - 1)(t_f - t_i)$$

$$\sigma^2 = k^2 4[p(1 - p)](t_f - t_i) \quad \text{Eq. 19}$$


$$\sigma^2 = k^2 4[p(1 - p)](t_f - t_i) \quad \text{Eq. 20}$$

It is now possible to consider another factor that may drive statistical drift of the size distribution in these models: differential rates of growth and shrinking, denoted  $k_g$  and  $k_s$ , respectively. The influence of differential rates of growth upon  $\mu$  and  $\sigma$  is more readily understood if  $k_g$  and  $k_s$  are re-framed into the drift of the size-frequency distribution and oscillations about the centre of the distribution (Fig. 6). The oscillatory component is  $k_{av} = (k_g + k_s)/2$ , the average rate with respect to the centre of the distribution, and the drift component is  $k_{net} = (k_g - k_s)/2$ , the imbalance in rates. The oscillations behave exactly as they do for a stationary distribution; so  $k$  becomes  $k_{av}$  in the equations above. Drift induced this way purely displaces the distribution, and so only affects  $\mu$ , adding a term so as to cause it to increase at a constant rate with time. Eq. 21 and Eq. 22 therefore describe a model [M7] combining Brownian motion randomness in growth with an exponential growth rate that includes the potential for overall growth of the population to be driven by both different probabilities and/or rates of growth and shrinkage; we term M7 the ‘*stochastic instability*’ (SI) model. With shrinking forbidden ( $k_s = 0$ ) and conceptualised in terms of discrete events, this simplifies to the model of Fowler et al. [22], which dealt with random uni-directional equally sized steps at a single rate creating growth.

$$\mu = \ln(H_i) + t[k_{\text{net}} + (2p - 1)k_{\text{av}}] \quad 465$$

$$\sigma^2 = k_{\text{av}}^2 4[p(1 - p)]t \quad 466 \quad \text{Eq. 21}$$

$$467 \quad \text{Eq. 22}$$



**Fig. 6 Illustration of how, conceptually, unequal rates of growth and shrinking may be decomposed into components.** The components represent: i), oscillation around the centre of a distribution of the logarithm of sizes; and ii), drift of the distribution.

Values for  $\mu$  and  $\sigma$  of the SI model [M7] may readily be estimated (see Appendix B) directly from mapped bedform sizes (e.g., Fig. 1). Through Eq. 21 and Eq. 22 the SI model therefore predicts trajectories of characteristics of the observed size distribution ( $\mu_{\text{obs}}$  and  $\sigma_{\text{obs}}$ ) through time; specifically  $\mu_{\text{obs}}$  is expected to be proportional to the square of  $\sigma_{\text{obs}}$ .

It is also possible to make predictions about the size differences (e.g.,  $\Delta H$ ) of flow-sets of bedforms across an observational window (i.e., at  $t_1$  and  $t_2$ ). First, all bedforms should be active and change size, and there should be a mixture of shrinking and growing. Secondly, in spite of the scatter caused by randomness,  $\Delta H$  should relate to  $H$  (Eq. 4). Thirdly, by the definition of a diffusive Wiener process  $t_N$  in any time period is normally distributed, and thus the distribution of the differences in height  $\Delta H$  should be log-normal. Furthermore, since the time difference is known, parameters of the SI [M7] model (i.e.,  $p$  or  $k_{\text{net}}$ ,  $k_{\text{av}}$ , total duration of growth period) may be uniquely constrained (Table 2).

**Table 2: Table of testable predictions for the WT [M10] and SI [M7] models.**

|   | Characteristic                                      | Expectation: WT model [M10]                                                                                                                                                           | Expectation: SI model [M7]                                                                                                                                                                                                             | Test/Investigative method                                                                                                         |
|---|-----------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------|
| 1 | Size-frequency distribution                         | Gamma; through time or across $\Delta t$ . $\beta$ constant; $\alpha \propto t$                                                                                                       | Log-normal through time or across $\Delta t$ . $\mu \propto \sigma^2 \propto t$                                                                                                                                                        | Repeat survey under active ice, or plot palaeo-forms from multiple flow sets (e.g., $\mu_{\text{obs}}$ vs $\sigma_{\text{obs}}$ ) |
| 2 | Spatial pattern of ice flow variables or conditions | Poisson fluctuations in time, at least at a bedform scale                                                                                                                             | Constantly fluctuating, at least at the spatio-temporal scale of bedform genesis                                                                                                                                                       | Estimate basal ice conditions using geophysics or invert for them from satellite observations of the ice surface [e.g., 6]        |
| 3 | Fraction shrinking vs growing                       | All active forms grow (i.e., $\Delta H$ is +ve)                                                                                                                                       | All active. $\Delta H$ a mixture of growing and shrinking; fraction $p$ growing.                                                                                                                                                       | Repeat survey under active ice; e.g., repeat [13]                                                                                 |
| 4 | Growth rate                                         | Constant. With $\Delta t$ known, $\Delta \alpha$ and $\Delta \beta$ are constrained and so are $\lambda$ and $k$ (Eq. 25, Eq. 26), so overall time to create flow set also deducible. | Exponential, i.e., proportional to $H$ . If $\Delta t$ known, $\Delta \mu$ and $\Delta \sigma$ and so $p$ or $k_{\text{net}}$ and $k_{\text{av}}$ are constrained (Eq. 21, Eq. 22), so overall time to create flow set also deducible. | Repeat survey under active ice.                                                                                                   |
| 5 | Fraction unchanged                                  | $>0$ for small $\Delta t$                                                                                                                                                             | Small; depends on definition of change                                                                                                                                                                                                 | Repeat survey under active ice.                                                                                                   |

## 5.2. Waiting time randomness [M8 to M11]

In contrast to Brownian motion randomness, there is another well-established type of temporal randomness called Poisson randomness [e.g., 94]. This is investigated in models M8 to M11.

In ‘Poisson’ randomness, the gaps between events that occur randomly at a given rate ( $\lambda$ , number per unit time) are distributed according to the exponential or ‘*waiting time*’ distribution [e.g., 97, p39-40]. This distribution is, for instance, used to model the times between shoppers arriving at a supermarket checkout. So, if the arrival or ‘event’ is the change in state (i.e., growth to inactivity) of a continuous process [cf. 91] it also describes inter-event periods in which bedforms may grow (Fig. 4b). Thus, if only a single episode of growth (e.g., the last) is preserved, net time spent growing ( $t_N$ ) is distributed according to an exponential distribution (Eq. 23).

501

$$f_{T_N}(t_N) = \lambda e^{-\lambda t_N}, t_N > 0$$

Eq. 23

As in Section 5.1, this is formulated in terms of time spent growing so that any desired growth rate law can be readily applied to determine distributions for  $H_f$ . The distributions of  $H_f$  that are generated by taking  $t_N$  as a random variable can be deduced by transformations of random variables as above [e.g., Ch 5 of 93].

Consider first model M8, in which growth is constant with time (Eq. 1). With  $t_N$  as above, an exponential distribution of heights results (Eq. 24). This, however, is not so for exponential growth (Eq. 14) in model M9. This produces a distribution that is not exponential, log-normal or Gamma. M8 predicts that the exponent of the tail of the observed pdf of final heights ( $H_f$ ) is  $\lambda/k$  as in Eq. 24, where growth rate ( $k$ ) is from Eq. 13. This exponent is readily estimated from mapped sizes [19], and is not expected to progress with time. It is predicted to be set by, vary in equilibrium with, and therefore reflect formative (i.e. ice or water) flow conditions.

$$f_{H_f}(h_f) = \frac{\lambda}{k} e^{-\lambda(\frac{h_f - H_i}{k})}, h_f > H_i \quad \text{Eq. 24}$$

However, instead of being in equilibrium with flow, glacial bedforms may be in a transient state with respect to flow. This is incorporated within models M10 and M11. If bedforms are created by a number ( $n_b$ ), on average, of building episodes then  $t_N$  is the sum of  $n_b$  exponential distributions; this is a two-parameter Gamma distribution denoted  $t_N \sim \Gamma(\alpha, \beta)$  [84]. The Poisson rate ( $\lambda$ ) as defined above is now standardly denoted  $\beta$  and is the ‘rate parameter’ of the Gamma distribution. The shape parameter of the Gamma distribution ( $\alpha$ ) is simply equal to  $n_b$  [e.g., 97, p292]. On average in M10 and M11 the number of building episodes is a multiplication of the rate at which they occur and the time that has elapsed, namely  $n_b = 0.5\lambda t$ , which is illustrated in Fig. 4b. The factor of 0.5 arises because two switches (‘on’ and ‘off’) are needed for each growth period.

The distributions of  $H_f$  that are generated in these Poisson multi-event models [M10 and M11] can be deduced by taking  $t_N$  as a Gamma distributed random variable, using growth

rates in equations Eq. 13 and Eq. 14, and as in previous sections then using transformations of random variables (i.e., Eq. 29). M10 has constant growth (Eq. 13), we term it the ‘*waiting time*’ (WT) model, and a Gamma distribution of heights results. This is not so for exponential growth (Eq. 14) upon which model M11 is based, which produces size distributions that are neither log-normal or Gamma.

The parameters of the WT [M10] model (i.e.,  $\lambda$ ,  $k$ , and  $t$ ) may be constrained from the rate ( $\beta$ ) and shape ( $\alpha$ ) parameters of the final height distributions ( $H_f$ ). They are related as in Eq. 25 and Eq. 26. Observed values are denoted  $\beta_{\text{obs}}$  and  $\alpha_{\text{obs}}$ , are readily estimated (e.g., figure 1 of [19]), and are predicted to be constant and increase linearly with time respectively.

$$\beta = \lambda/k \quad \text{Eq. 25}$$

$$\alpha = n_b = 0.5\lambda(t_f - t_i) \quad \text{Eq. 26}$$

It is possible to make predictions about the size differences (e.g.,  $\Delta H$ ) expected across a time window (i.e., at  $t_1$  and  $t_2$ ). First, all bedforms that have changed should have grown, and a fraction should not have changed if the number of building events ( $n_b = \alpha$ ) is small. Secondly, growth should be at a constant rate and  $\Delta H$  should not correlate strongly with  $H$  (Eq. 1). Thirdly, the ‘memoryless’ nature of the Poisson process dictates that  $\Delta H$  should be a Gamma distribution. Furthermore, since the time difference is known, the rate constant of bedform growth ( $\lambda$ ) could then be estimated uniquely through the two observations of  $\alpha$  (i.e.,  $\Delta\alpha_{\text{obs}} = \alpha_2 - \alpha_1 = 0.5\lambda\Delta t$ ). Then, growth rate ( $k$ ) could be calculated through either observation of  $\beta$  (see Table 2).

## 6. Results

The right hand column of Table 1 lists which models produce size-frequency distributions that have been argued to reasonably approximate mapped observations (i.e., log-normal[22,35,36], gamma, or exponential above mode[19]). Fig. 1 shows a direct comparison, illustrating how well each of these three alternatives fit the data: solid line is an exponential



distribution, generated by model M8; dashed line is a log-normal distribution generated by M7, the Stochastic Instability (SI) model; dotted line is a gamma distribution generated by M10 the Waiting Time (WT) model. Other models, however, can fit. By invoking substantial *ad hoc* assumptions (see Appendix A), some models that do not involve stochasticity in growth through time [M3a, M4a, M5a] can also replicate size-frequency observations. Fig. 2, and Figs 8 to 10 in Appendix A, also show some of the shapes generated by the other models. It is important to note that fitting statistical distributions as in Fig. 1 in itself leads to parameters (e.g.,  $\mu$  and  $\rho$ , or  $\phi$  and  $\lambda$ ) that are only descriptive empirical quantities; it is the statistical bedform growth models that relate the parameters to key aspects of the physical process: antecedent topography, growth rate (e.g., exponential), and the timing of growth.

## 7. Discussion

To gain additional insight into the plausibility of conceptual models of the growth of subglacial bedforms, this paper takes well-established statistical behaviours (e.g., types of temporal randomness) and integrates them with plausible growth rate behaviours [e.g., 20] to explore which combine to produce reasonable approximations of the observed size-frequency distribution of subglacial bedforms (i.e., exponential, Gamma, or log-normal [e.g., 19,22]). Exactly as any model (e.g., numerical ice sheet models) these contain approximations and assumptions, but are constructed to capture key aspects of the physical processes in order that these might be evaluated by comparing modelled outputs to observations. In 7.1, the statistical models [M1-M11] are evaluated in terms of their ability to explain i) the size-frequency observations whilst invoking the least number of *ad hoc* or arbitrary assumptions, ii) their internal consistency, and iii) their ability to explain all other relevant observations (e.g., geophysics). The implications of the favoured model are then discussed (section 7.2), followed by some suggestions for future work (section 7.3).


### 7.1 Evaluation of the models

580 The simplest models created [M1-5] do not involve stochasticity in growth through time.  
 581 For any of these (see Table 1) to replicate size-frequency observations (Fig. 1) they require  
 582 substantial *ad hoc* assumptions or special pleading, discussed in Appendix A. This we  
 583 interpret as making these models, as constructed, less plausible and giving some weight to the  
 584 view that neither ‘classic’ deterministic growth nor antecedent bedform-scale topography are  
 585 sufficient to explain bedform sizes. It should be noted, however, that the failure of one  
 586 particular modelling realisation of an envisaged process rarely excludes that process.

587 Models M6 to M11 follow up on the conceptual model of [19] in that they are based on  
 588 variations in growth through time. Constructions M6 and M9 do not match the size-frequency  
 589 observations (Table 1) and they can be ruled out. M8 can reproduce the exponential tail, but  
 590 to allow it to fit the data fully it must either invoke selective post-formational degradation or  
 591 an argument that observational data have missed most small bedforms in order to create the  
 592 roll-over. This is debatable; first, even the ~25% recovery rate affecting small drumlins is  
 593 insufficient to wholly explain the roll-over in the UK data [31,102], and second the very many  
 594 small forms expected of an exponential distribution are mapped in high-resolution data of  
 595 neither previously glaciated [e.g., 103] nor recently uncovered [40] drumlin fields. In contrast  
 596 to M8, both types of temporal randomness, when combined with appropriate growth rates into  
 597 the SI and WT models (i.e., in M7 and M10, but not M6 or M11), fit the widespread palaeo-  
 598 bedform size data. Neither Poission nor Brownian Motion randomness in growth have yet  
 599 been specifically identified under active ice, but they have been observed commonly in  
 600 natural processes including bedform evolution [25–28,30,57,80,92,96], and so are supported  
 601 by analogy. This, we argue, makes their introduction significantly less *ad hoc* than the  
 602 arbitrary assumption of convenient statistical distributions in M3a to M5a. Note, for instance,  
 603 that the temporal variation that distributes  $t_N$  in the SI model [M7] intrinsically creates the  
 604 Gaussian distribution arbitrarily invoked by M5a.

605 Significantly, and in their favour, models M7 (‘stochastic instability’: SI) and M10  
 606 (‘waiting time’: WT) also explain other independent observations of bedforms without any

607 further *ad hoc* additions. First, probabilistic growth decouples initial and final sizes, allowing  
608 the intervening physical process to dominate the characteristics of the ultimate size-frequency  
609 distribution; that is, illustratively, the randomness in growth shown in Fig. 7 dictates the size-  
610 distribution, not the initial size. This offers an explanation for the observation that drumlins  
611 with their typical size-distribution can originate irrespective of differences in environment  
612 (e.g., till/bedrock lithology) [42,43]. Secondly, the observed structure (e.g., internal  
613 stratigraphy [e.g., 12,40]), the variety of composition [e.g., 42,43], and the substantial (e.g.,  
614  $\pm 50\%$ ) scatter in the sizes and elongations commonly seen for proximal palaeo-forms within a  
615 flow-set [e.g., 16,39,45,104], might be expected to result from randomness and fluctuations  
616 in characteristics of the ice-sediment-water system in space and time. By their design, the WT  
617 and SI models are also consistent with the geophysical, remotely sensed, and  
618 sedimentological evidence for spatio-temporal variability in ice flow velocity and the bed  
619 beneath ice sheets, which was outlined in sections 3.1 and 3.2. Thus, the widespread dataset  
620 of palaeo-bedform sizes points towards a view where ice-water-sediment dynamics (i.e.,  
621 change through time) likely has a fundamentally random element that physics-based models  
622 of bedform genesis could usefully incorporate; to date, some models have been seeded with  
623 initial random height perturbations [48,79], but what if any temporal randomness to emerge  
624 from this has not been explicitly examined. Fowler et al. [22] demonstrated that a statistical  
625 model can reconcile observations with the hypothesis of Hillier et al. [19], but the variety of  
626 statistical models considered here allows us for the first time to distinguish process dynamics  
627 (i.e., randomness through time) as the most plausible origin for the necessary variability out of  
628 the main candidates.



**Fig. 7 Evolution of bedforms including randomness through time.** The evolution of sizes of ten illustrative bedforms including randomness in their growth through time (grey lines). These differ from a deterministic path (black line). For a sufficiently large number of bedforms, the average properties (e.g., mean size) of a flow set closely approximate the deterministic path. Bedforms are ‘born’, last pass a threshold minimum observable height (e.g., 1 unit, dashed line), at different times.

It is possible to argue that one type of bedform-scale dynamics is more likely, i.e., differentiate between the SI [M7] and WT [M10] models. First, by visual inspection the log-normal shape produced by the SI model arguably fits the size-frequency data than the gamma distribution of the WT model, especially for  $L$  and  $W$ , and for small sizes (see Fig. 1). Secondly, it allows bedforms to shrink as seems probable from the geophysical observations [11,12], which the WT model does not. Thirdly, the SI and WT models may also be evaluated through their internal consistency between observations for the three dimensions  $H$ ,  $W$ , and  $L$ . Taking the simplest assumption that all dimensions change size together (i.e.,  $t$  and  $p$  are the same), Eq. 22 can be used to constrain relative growth rates (e.g.,  $k_{avH}/k_{avW}$ ) for the dimensions within the SI model (Eq. 27). Values for  $\sigma$  calculated for mapped UK drumlin data given in Fig. 1 then indicate that increasing  $H$  is the primary mode in their genesis, namely its growth rate constant is greatest ( $k_{avH} > k_{avL} > k_{avW}$ ). This is plausible. In contrast, using Eq. 26,  $\alpha$  values for the WT model [M10] imply a different number of growth episodes for each dimension. This is less easily explicable. Thus, with these factors taken together, we choose to favour the SI model over the WT model.

$$\frac{\sigma_L}{\sigma_W} = \frac{k_{avL}}{k_{avW}}, \quad \frac{\sigma_H}{\sigma_W} = \frac{k_{avH}}{k_{avW}}, \quad \frac{\sigma_L}{\sigma_H} = \frac{k_{avL}}{k_{avH}} \quad \text{Eq. 27}$$

Alternatively, stochasticity in the ice-sediment-water system may differ from the Brownian motion of our SI model, but with exponential growth still produce log-normal size-frequency distributions because of the central limit theorem (CLT) [22]. Fowler et al. [22] interpret this as favouring growth through discrete 'events' of constant size, but the CLT has other interpretations [e.g., 105:p88,106:p266], so this is not necessarily required. For instance, if growth of each bedform is governed by discrete 'events' of random size, selected from any frequency distribution, the CLT predicts a log-normal distribution of sizes in a flow set. Similarly, if bedforms grow by many growth periods of a random duration selected from any frequency distribution, the CLT dictates that effective  $t_N$  will be Gaussian as required. However, even given this, the SI model is still likely to be a useful *empirical approximation*. If the factors dictating bedform-scale randomness (e.g., supra-glacial lake drainage patterns) relate to broader ice-sediment-water conditions then parameters fitted as for the SI model (i.e.,  $\mu$ ,  $\sigma$ ) will still provide a useful statistical link between observations at the flow-set level and theory such as in numerical ice flow models (e.g., by plotting spatial distributions).

## 7.2 Implications of the SI approximation

The SI model, if it is to be accepted as most likely, has a number of implications. Bedforms are expected to change size randomly through time in a manner approximating Brownian




Fig. 7). The quantitative, observable corollaries of this are listed in Table 2. A number of

points, however, need some further explanation.

First, the SI model implies that it is not necessary to invoke a lower ‘physical threshold’ on drumlin length or width [16] or an upper limit for  $H$  a quenching (a.k.a. ‘capping’) mechanism to limit their upper ‘critical size’ [e.g., 20,77,78,107]. In the SI model very small sizes are simply less likely and no lower threshold is needed. As an alternative explanation for the absence of extremely large bedforms, the SI model and its simpler variant [i.e., 22] must invoke growth that is ‘transient’, namely that it occurs within a time window of limited duration. Simply, insufficient time has passed for very large forms to be created. Observations of active bedforms do not yet indicate which means of limiting the largest sizes is most plausible, but several mechanisms can be imagined that allow growth periods forming flow sets to be of limited duration. In a steady-state view, meso-scale patches of bedforms could be periodically flattened by conditions adverse to the existence of bedforms. Alternatively, favourable patches may only occur transiently [e.g., 39] or time-transgressively [e.g., 38] as ice sheets melt and retreat. However, to explain bedform prevalence, these mechanisms must commonly occur. Size-frequency observations give two tentative indications that a time limitation (e.g., SI model) affects glacial bedforms rather than a physical cap in an equilibrium model [e.g., 78]. The first indication is that fluvial bedforms measured at

equilibrium with flow do not have a log-normal distribution, but one that peaks at larger sizes [Fig. 6a of 26] as if sizes were tending to bunch below some fuzzy threshold. The second indication is that if glacial bedforms were to grow and then to ‘freeze’ [78] at a sharp upper limit a peak in frequencies would be expected, but this is not observed in Fig. 1c (i.e., at 34 m).

Secondly, assuming all dimensions change size together (i.e.,  $t$  and  $p$  are the same), relative growth rates estimated from UK observations (Fig. 1, Eq. 27) (i.e.,  $k_{avL} > k_{avW}$ ) indicate that drumlins elongate as they grow [e.g., 16,31]. Note that no relationship between the dimensions was placed into the SI model that might have prescribed this observation. Perhaps they continue into mega-scale glacial lineations (MSGSL) as part of a genetically-linked bedform continuum [cf. 108,109], where  $H$  and  $W$  are in equilibrium restricted by stochastic interactions with ice and neighbouring bedforms whilst elongation continues.

Thirdly, Fowler et al. [22] put forward an explanation to demonstrate that size observations do not necessarily falsify the exponential growth hypothesised in the physically-based till ‘instability models’ of bedform genesis [e.g., 20]. Here, a variety of different explanations are considered, and exponential growth still features in the one that is apparently most plausible. Thus, through this comparison, the SI model strengthens the tentative observational support for exponential bedform growth (i.e. by linear instability). On the other hand, from two-parameter fits to observed data collated in a small number of distributions (e.g., Fig. 1) it is not possible to distinguish between existing linear instability mechanisms, namely till or heat-flux [e.g., 20,33]. Future work plotting the spatial distribution of parameters ( $\mu$ ,  $\sigma$ ) of mapped palaeo-bedforms against numerically modelled predictions of growth rate ( $k$ ) for each mechanism for a past ice sheet could, however, distinguish them. Other possible tests and applications of the SI model are considered below.

### 7.3 Future Work: Testing and applying the SI model

The SI model [M7], if correct, suggests tentative analytical links between parameters fitted to observed size-frequency distributions and ice sheet properties, such as ice velocity; the SI

model links size observations ( $\mu$ ,  $\sigma$ ) to growth rate  $k$  (Eq. 21 and Eq. 22), which relates to physical parameters [e.g., 33]. Eq. 52 of Fowler [110], for instance, related  $k$  to  $(AN/2\eta)^{1/2}$  within which  $A$  is illustratively proportional to ice velocity. Similarly, Shoemaker [56] related  $k$  to subglacial flood water velocity to a power  $\frac{16}{3}$ . Thus, predicted relationships (e.g.,  $k \propto \sqrt{v}$ ) can contribute to geomorphological debates such as the interpretation of  $L$  in terms of  $t$  or  $v$  [e.g., 3]. Admittedly, the problem is under-constrained since there are three variables ( $p$  or  $k_{\text{net}}$ ,  $k_{\text{av}}$ , and  $t$ ) and two observables ( $\mu$ ,  $\sigma$ ). If, however, more can be learnt about one of these through direct observation or experimentation (e.g.,  $p$ ) the other two (e.g.,  $t$  or  $k$ ) could be determined remotely from a single morphometric analysis.

The SI model makes quantitative predictions that are distinctively different from the WT model or deterministic ones, as detailed in Table 2. This makes it testable and falsifiable by observations from modern subglacial environments. The predictions are, for example, testable by repeating at  $t_2$  a past (i.e., at  $t_1$ ) geophysical survey under active ice [i.e., 13]. In addition, plots of size-frequency parameters obtained for a number of observed flow sets are diagnostic of different models (see Section 5); for instance, in the SI model  $\mu \propto \sigma^2$ , so plots of  $\mu$  against  $\sigma^2$  will display linear trends if  $t$  varies whilst the other variables are held constant. Plotting spatial variations in parameters could also be an additional constraint upon physics-based models of bedform genesis. Illustratively, consider a numerical model used to estimate ice flow in a past ice sheet [e.g., 111], a physics-based model of bedform genesis [e.g., 33], and a hypothesised set of conditions (e.g., based on basal shear stress) for drumlin formation. Then, the modelled ice-sheet conditions set  $t$  for flow-sets geomorphologically mapped for that ice sheet, and in conjunction with the model of bedform genesis they also set a numerical prediction for  $k$ . Furthermore, since  $t$  is constrained in the context of this test,  $k$  and  $p$  can be determined for the mapped flow sets by using a statistical model (see above). Thus, through the spatial distribution of  $k$ , a way exists to quantitatively compare models and observations. Patterns in  $k$  could either be of absolute or relative values, and  $k$  and  $p$  may relate to properties of ice flow (e.g.,  $v$ ) or postulated floods depending upon the drumlin formation model



selected. In particular, the ability or not to correctly predict the distribution and properties of flow sets may help to further constrain which ice sheet models, or members of an ensemble of potential realisations, is most valid.

Since we do not attempt to develop all possible models here, the wider point is that statistical modelling provides a tool to develop and falsify conceptual models of bedform growth. The same is true for other bedforms where measurement of key processes is challenging (e.g., in-situ on barchan dunes) and where time-series of digital elevation models are becoming available but statistical work is limited [e.g., 18]. With respect to fluvial environments, developing our analytical work could create statistical distributions reflecting underlying mechanics, improving upon existing distributions as descriptors [e.g., 26] and allowing more to be extracted from field observations.

## 6. Conclusions

The emergence and growth of subglacial bedforms is difficult to observe, significantly limiting our ability to accurately parameterise basal processes beneath ice sheets. In this paper, a novel approach has been taken, developing new probabilistic growth models and comparing their predictions with observed distributions of palaeo-bedform sizes. The variety of explanations both permits a number of models to be discounted and the relative plausibility of the rest to be assessed for the first time. The ‘*stochastic instability*’ (SI) model, modified from Fowler et al. [22] and extended to encompass bedforms shrinking, is argued to provide the best fit to observations. Not only does it fit the size observations [22], but it appears to do so with fewest *ad hoc* assumptions whilst being internally self-consistent between metrics (e.g., height and width) and in accord with other observations (e.g., geophysical). Thus, our analysis strengthens a view [19,22] where the ice-sediment-water dynamics and sediment flux have significant elements of randomness in space and time (i.e., not continuous or monotonic) and cause both erosion and deposition. This view is developed to explicitly argue that (i) flow-related processes at the ice-bed interface rather than initial bedform-scale topography

773 govern bedform sizes and (ii) drumlins elongate with time. Furthermore, parameters of  
774 mapped size-frequency distributions are explicitly linked with ones related to flow (i.e. ice  
775 and water) for the first time, accompanied by an illustration of an avenue for how this may be  
776 used to improve calibration of basal conditions in numerical ice sheet models and achieve a  
777 better understanding of conditions at the base of ice sheets. Lastly, we demonstrate that it is  
778 possible to provide testable, distinctive predictions that will allow models to be distinguished  
779 using a hypothesised repeat geophysical survey of bedforms under active ice. Note that none  
780 of the work presented here precludes or conflicts with observations of structured spatial  
781 patterning in the bedforms.

782


## 783 **Appendix A: Preliminary exploration**

784       Following the trajectory of work that developed stochastic sub-aerial landscape evolution  
 785 models to explain topography's typical fractal statistics [112], this appendix formalises  
 786 statistically for the first time simple models representing the prevailing 'classic' view that  
 787 bedform growth through time is not random, which has not yet been undertaken for subglacial  
 788 bedforms. In these simpler models, elements of the potential spectrum of randomness within  
 789 the proposed meso-scale patches are, effectively, turned off.

790       The first models [M1-3] represent the more plausible realisations of the 'classical' view  
 791 where bedform growth through time is not random. M1 considers the simplest, entirely  
 792 deterministic, case. It is possible that the bedform-scale topography prior to bedform creation  
 793 is not planar, so models M2 and M3 include variability in initial bedform height. It has also  
 794 been proposed that bedforms are not 'born' at the same time [cf. 11,113], so models M4 and  
 795 M5 assess the possibility that each bedform could start to grow at a different time. The models  
 796 are described then evaluated.

### 797 **A.1. Entirely deterministic growth [M1]**

798       Model M1 considers multiple independent bedforms all of a single initial height ( $H_i$ )  
 799 growing according to any given deterministic mechanism; the 'classical' view that has yet to  
 800 be explicitly tested. The bedforms will all reach the same final height ( $H_f$ ) as each other after  
 801 any time has elapsed (i.e.,  $t_f - t_i$ ), whatever their growth rate (Fig. 8). This model starts with a  
 802 Dirac delta function as the pdf (probability density function) of  $H_i$  and produces the same pdf  
 803 of  $H_f$  at a later instant in time  $t_f$ , namely a single vertical spike on plots such as Fig. 2 or Fig.  
 804 8.




**Fig. 8 Probability density functions (pdfs) for the simplest model [M1].** In this model drumlins have a single initial height  $H_i$ , then grow deterministically through time.

### A.2. Variable initial topography [M2 and M3]

Models M2 and M3 are designed to give insight into whether or not the observed final size-frequency distribution may simply arise as a result of an inherited distribution of initial sizes, without recourse to stochastic behaviour during growth. These models are stochastic in the initial conditions only; that is, the initial condition of Eq. 2 is modelled as a random variable following a prescribed pdf that reflects a chosen initial size distribution.

Proto-bedforms of initial height  $H_i$  follow a uniform distribution, that is they are equally distributed across a range of heights between  $a$  and  $b$  (Eq. 28), which is the width of the grey boxes on Fig. 9, and grow deterministically.

$$f_{H_i}(h_i) = \begin{cases} \frac{1}{b-a}, & \text{for } a < h_i < b \\ 0, & \text{elsewhere} \end{cases} \quad \text{Eq. 28}$$



**Fig. 9 Pdfs for models with deterministic growth and variable initial topography a)** linear growth [M2] **b)** exponential growth [M3]. Initial  $H$  distribution  $H_i$  (grey, dashed line) changes to the final one  $H_f$  (black outline) as time progresses. Dotted lines are an arbitrary function. Cases shown are where smallest  $H_i$  is zero;  $a = 0$ .

So defined,  $H_i$  is a random variable; thus, since  $H_f$  in Eq. 3 and Eq. 5 is a function of  $H_i$ , it is also a random variable whose distribution can be determined. Determining the pdf of  $H_f$  given the pdf of  $H_i$  is a relatively straightforward task. To this aim, the standard transformation relationship

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right| \quad \text{Eq. 29}$$

relating random variables  $y$  and  $x$  is invoked assuming a relationship of the form  $y = g(x)$  [e.g., Ch 5 of 93].

If growth is linear with time (Eq. 1) [M2], the shape of the initial distribution is not altered (Eq. 30) and it moves right as illustrated in Fig. 9a. So, if any non-trivial growth (e.g., 4 m) has occurred, it is not possible to construct a pdf for  $H_i$  that still contains low amplitude bedforms; for example, even the smallest initial height of 0 m would have grown to 4 m. For mapped size data the mode ( $\phi_{\text{obs}}$ ) would increase linearly with time, but the exponent of the right-hand tail ( $\lambda_{\text{obs}}$ ) [19] would stay constant.

$$f_{H_f}(h_f) = \begin{cases} \frac{1}{b-a}, & \text{for } a + k(t_f - t_i) < h_f < b + k(t_f - t_i) \\ 0, & \text{elsewhere} \end{cases} \quad \text{Eq. 30}$$

If growth is caused by linear instability [M3] (i.e., is exponential as in Eq. 4) then the distribution elongates (Eq. 31, Fig. 9b) but does not alter the relative abundances of different bedform sizes (e.g., 5th, 50th and 95th percentiles of  $H$ ). Indeed, the pdf can be imagined as being drawn on a sheet of elastic material so that, even if it is any arbitrary function (dotted lines), it will be elongated but not otherwise distorted. Thus, to end up with an approximately log-normal distribution as observed for bedforms (e.g., Fig. 1), a landscape must start with a log-normal distribution; this *ad hoc* modification of M3 is denoted M3a. For mapped size data M3a would have both  $\phi_{\text{obs}}$  and  $1/\lambda_{\text{obs}}$  increasing linearly proportional to each other and with the duration of the bedform building episode, and this would happen along a trajectory set by the shape of the initial distribution.

$$f_{H_f}(h_f) = \begin{cases} \frac{1}{(b-a)e^{k(t_f-t_i)}}, & \text{for } ae^{k(t_f-t_i)} < h_f < be^{k(t_f-t_i)} \\ 0, & \text{elsewhere} \end{cases} \quad \text{Eq. 31}$$

### A.3. Variable initiation times [M4 and M5]

Models M4 and M5 formalise the glaciological hypothesis in which bedforms are not ‘born’ at the same time and therefore, at any point in time, will have been growing for different durations [11,113]. Proto-bedforms of an initial (constant) size  $H_i$  start growing at times distributed according to a uniform distribution from an earliest time defined as  $c$ ; i.e., a constant number are created per unit time as the building of the flow set progresses. All continue growing until a final, constant time ( $t_f$ ). The time at which bedforms’ growth starts,  $t_i$ , is now a random variable (Eq. 32) making final height ( $H_f$ ) also a random variable since it is a function of  $t_i$ . The pdf of  $H_f$  can be determined similarly to the previous section by resorting to the transformation relationship of Eq. 29.


867

$$f_{T_i}(t_i) = \begin{cases} \frac{1}{t_f - c}, & \text{for } c < t_i < t_f \\ 0, & \text{elsewhere} \end{cases} \quad \text{Eq. 32}$$

869

870

871 If growth is linear with time (Eq. 1) [M4], then a uniform distribution of final heights is



872 produced (

873 Fig. 10a, Eq. 33). In general, *ad hoc* manipulation of the form of the pdf of  $t_i$  will be

874 directly reflected in the output form of  $H_f$ . A linearly increasing production rate (number per

875 unit time), for instance, would produce a linearly decreasing frequency with increasing  $H_f$

876 because the larger number of recently produced forms have not yet had time to grow. Thus, an

877 approximately Gamma distribution (e.g., Fig. 1), for instance, could be created by a


878 production rate that started slowly, built approximately exponentially to a peak and then died

879 rapidly before  $t_f$ ; this variant is denoted M4a. If interrupted at any point before the distribution

880 was fully formed, the distribution would have its left side missing as this part would not yet

881 have been created. In terms of mapped size data,  $\phi_{\text{obs}}$  would remain at  $\sim 0$  until the roll-over

882 was created, and  $1/\lambda_{\text{obs}}$  would remain constant if the right hand tail were well-approximated  
 883 by an exponential distribution.




884 **Fig. 10 Pdfs for models with deterministic growth where bedforms have constant**  
 885 **initial heights, but a uniform distribution of initiation times (i.e., initiation rate is**  
 886 **constant through time) a) linear growth [M4] b) exponential growth [M5]. Initial**  
 887 **distribution (grey, dashed line) changes to the final one (black outline).**  
 888  
 889

$$f_{H_f}(h_f) = \begin{cases} \frac{1}{k(t_f - c)}, & \text{for } H_i < h_f < H_i + k(t_f - c) \\ 0, & \text{elsewhere} \end{cases} \quad \text{Eq. 33}$$



894 If growth is exponential (Eq. 4) [M5], the frequency of remnant forms is not exponential,  
 895 but is inversely proportional to final height (Eq. 34,



896  
 897 Fig. 10). This is verifiable intuitively since frequency in any height band is less the faster  
 898 bedforms pass through it; specifically, bedform frequency is inversely proportional to their  
 899 growth rate (i.e.,  $1/kH$ , Eq. 4). In order to replicate an approximately log-normal distribution  
 900 of  $H_f$  (e.g., Fig. 1) with exponential growth,  $t_i$  must have a roughly Gaussian (i.e., normal)  
 901 distribution [M5a]; a log-normal distribution is defined as that of a random variable whose  
 902 logarithm is normally distributed, and Eq. 4 can be written to give the logarithm of  $H_f$  as  
 903  $\log(H_f) = \log(H_i) + k(t_i - c)$  where everything on the right hand side is constant here except  
 904  $t_i$  which is a normal distribution. This can be verified by appropriate transformations of the  
 905 random variables [e.g., Ch 5 of 93]. Giving  $t_i$  a normal distribution would, strictly, allow it to  
 906 take values from  $-\infty$  to  $+\infty$ , and so to apply to a period of bedform creation ranging between  $c$   
 907 and  $t_f$  only *ad hoc* Gaussians with small values outside this range could be employed. For  
 908 mapped size data M5a predicts that  $1/\lambda_{\text{obs}}$  would increase linearly with time along a  
 909 trajectory set by the shape of the initial distribution, and  $\phi_{\text{obs}}$  would remain at  $\sim 0$  until the  
 910 roll-over was created, then increase exponentially. Note that the SI model [M7] gives a

mechanistic explanation for a Gaussian distribution of net growth durations rather than an *ad hoc* assumption of this in M5a.

$$f_{H_f}(h_f) = \begin{cases} \frac{1}{h_f H_i e^{k(t_f - c)}}, & \text{for } H_i < h_f < H_i e^{k(t_f - c)} \\ 0, & \text{elsewhere} \end{cases} \quad \text{Eq. 34}$$

#### A.4. Evaluation of models M1 to M5

With no randomness or variation [M1], the observations cannot be replicated. That is, no sharply spiked peaks are observed in size frequency distributions (Fig. 1), casting serious doubt upon an entirely deterministic model. Thus, M1 is rejected. M2 and M3 are based on variations in initial bedform sizes,  $H_i$ . Linear deterministic growth with uniformly distributed initial heights [M2] does not retain the small forms that are observed. Indeed, as explained above, there is no distribution of initial heights that can do so. Similarly, linearly unstable (i.e. exponential) deterministic growth [M3] does not intrinsically create an appropriate, exponentially tailed, size-frequency distribution. A progenitor landscape with log-normal  $H_i$  must be invoked to give the required log-normal  $H_f$  [M3a], but this *ad hoc* modification is somewhat questionable in a world where fractals (i.e., power-law distributions) dominate topography [e.g., 114]; even when suggesting that earlier progenitor log-normally sized forms may exist to be altered, the first set needs explaining. Thus, we provide the first observational constraint to indicate that something more appears to be needed than the ‘classic’ deterministic view of bedform growth and more obvious variants represented by models M1 to M3.

M4 and M5 are based on variations in growth initiation times,  $t_i$ . Linear deterministic growth with a uniform distribution of initiation times [M4] does not match the size-frequency distribution. *Ad hoc* manipulation [M4a] is therefore needed. However, M4a invokes, without supporting evidence or analogy, a ‘reflected’ log-normal distribution of frequency that starts slowly, builds approximately exponentially to a peak, and dies rapidly before  $t_f$ . Exponential

growth, as illustrated by a uniform distribution of initiation times [M5], does not intrinsically lead to an approximately Gamma or log-Normal distribution of bedform sizes that is observed. A Gaussian distribution (i.e.,  $t_i \sim N(\mu, \sigma)$ ) would explain the observations [M5a], but it must be arbitrarily invoked. Thus, if bedforms are ‘born’ at different times [see 11,113], it is demonstrated that a very specific pattern of ‘births’ is needed. Arguably, it would be preferable to have some process-related explanation for the required distribution of their initiation times.

## Appendix B: Parameter estimation

Descriptions of the calculation of the exponent ( $\lambda$ ) above a mode ( $\phi$ ) and parameters of a gamma distribution ( $\alpha_{\text{obs}}$ ,  $\beta_{\text{obs}}$ ) are given in Hillier et al. [19], which explicitly includes how counts from previously published size-frequency plots can be utilized. Fowler et al. [22] relays the standard formulae for a log-normal distribution where individual data are available ( $\mu_{\text{obs}}$ ,  $\sigma_{\text{obs}}$ ), and how this may be done for digitisations of previously published size-frequency plots is given below. Worked examples for all parameters and all the data sets used in this paper are provided in EXCEL sheets as Supporting Information.

Maximum likelihood estimation of log-normal distribution parameters ( $\mu$ ,  $\sigma$ ) using binned data, such as that digitised in Fig. 1, adapts standard formulae used to calculate  $\mu$  and  $\sigma$  for individual data in various areas of research [e.g., 22,115,116]. The mean,  $\bar{x}$ , and standard deviation,  $s_x$ , of the sample are calculated to estimate  $\mu$  and  $\sigma$ , respectively, using equations 35 and 36.  $n$  is the total number of data with counts,  $c_j$ , of bins at  $x_j$ .

$$\hat{\mu} = \bar{x} = \frac{1}{n} \sum c_j \ln(x_j) \quad \text{Eq. 35}$$

$$\hat{\sigma} = s_x = \sqrt{\frac{1}{n-1} \sum c_j [\ln(x_j) - \overline{\ln(x)}]^2} \quad \text{Eq. 36}$$

963

964 **Table of Notation**

965

| Symbol                                    | Quantity                                                                                                                                                              | Units                 |
|-------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| $i, f$                                    | Initial and final, e.g., referring to $H$ or $t$ .                                                                                                                    | n/a                   |
| $H, W, L$                                 | Height, width and length. Strictly, $H$ is bedform amplitude.                                                                                                         | m                     |
| $t$                                       | Time; $t_1$ and $t_2$ are earlier and later times respectively                                                                                                        | s                     |
| $t_N$                                     | Net time spent growing                                                                                                                                                | s                     |
| $t_g, t_s$                                | Time growing, shrinking                                                                                                                                               | s                     |
| $a, b, c$                                 | Constants                                                                                                                                                             | m, m, s               |
| $\alpha, \beta$                           | Parameters of the Gamma distribution – WT model [M10]; $\alpha_{\text{obs}}, \beta_{\text{obs}}$ are values of metrics estimated from observed size-frequency data.   | no units, $s^{-1}$    |
| $\mu, \sigma$                             | Parameters of the log-normal distribution – SI model [M7]; $\mu_{\text{obs}}, \sigma_{\text{obs}}$ are values of metrics estimated from observed size-frequency data. | no units              |
| $\lambda$                                 | Rate parameter for Poisson processes.                                                                                                                                 | $s^{-1}$              |
| $\lambda_{\text{obs}}, \phi_{\text{obs}}$ | Exponent and mode of size-frequency data, as approximated in Hillier et al. (2013).                                                                                   | $m^{-1}$ , m          |
| $k$                                       | Growth rate constant                                                                                                                                                  | $ms^{-1}$ or $s^{-1}$ |
| $n$                                       | Number of bedform observations.                                                                                                                                       | no units              |
| $k_g, k_s$                                | Growth rates of growth and shrinking, when differentiated; see text for relation to $k_{\text{av}}, k_{\text{net}}$ .                                                 | $s^{-1}$              |
| $n_b$                                     | Number of growth episodes – WT model [M10].                                                                                                                           | no units              |
| $j$                                       | Number of bedforms in a patch                                                                                                                                         | no units              |
| $p$                                       | Probability of growth                                                                                                                                                 | no units              |
| $\xi$                                     | Statistical drift – SI model [M7]                                                                                                                                     |                       |
| $v$                                       | Ice velocity                                                                                                                                                          | $ms^{-1}$             |
| $\tau$                                    | Basal shear stress                                                                                                                                                    | $Nm^{-2}$             |

966

967

968

## Acknowledgements

We thank the editors of PLOS ONE and three anonymous reviewers for their detailed and insightful reviews.

## References

1. Hart JK. Identifying fast ice flow from landform assemblages in the geological record: a discussion. *Ann Glaciol.* 1999;28: 59–67.
2. Kleman J, Hättestrand C. Frozen-bed Fennoscandian and Laurentide ice sheets during the Last Glacial Maximum. *Nature.* 1999;402: 63–66.
3. Stokes CR, Clark CD. Are long subglacial bedforms indicative of fast ice flow? *Boreas.* 2002;31: 239–249.
4. Schoof C. Basal perturbations under ice streams: form drag and surface expression. *J Glaciol.* 2002;48: 407–416.
5. Sergienko O V., Bindschadler RA, Vornberger PL, MacAyeal DR. Ice stream basal conditions from block-wise surface data inversion and simple regression models of ice stream flow: Application to Bindschadler Ice Stream. *J Geophys Res.* 2008; doi:10.1029/2008JF001004
6. Sergienko O V., Creyts TT, Hindmarsh RCA. Similarity of organized patterns in driving and basal stresses of Antarctic and Greenland ice sheets beneath extensive areas of basal sliding. *Geophys Res Lett.* 2014;41. doi:10.1002/2014GL059976
7. Siegert MJ, Taylor J, Payne AJ. Spectral roughness of subglacial topography and implications for former ice-sheet dynamics in East Antarctica. *Glob Planet Change.* 2005;45: 249–263. doi:doi:10.1016/j.gloplacha.2004.09.008
8. Vaughan DG, Smith AM, Nath PC, Le Meur E. Acoustic impedance and basal shear stress beneath four Antarctic ice streams. *Ann Glaciol.* 2003;36: 225–232.
9. Smith AM. Microearthquakes and subglacial conditions. *Geophys Res Lett.* 2006;33. doi:10.1029/2006GL028207
10. Murray T, Corr H, Forieri A, Smith AM. Contrasts in hydrology between regions of basal deformation and sliding beneath Rutherford ice stream, West Antarctica, mapped using radar and seismic data. *Geophys Res Lett.* 2008;35. doi:10.1029/2008GL033681
11. Smith AM, Murray T, Nicholls KW, Makinson K, Athalgerirdottir G, Behar A, et al. Rapid erosion and drumlin formation observed beneath a fast-flowing Antarctic ice stream. *Geology.* 2007;35: 127–130.
12. King EC, Woodward J, Smith AM. Seismic and radar observations of subglacial bed forms beneath the onset zone of Rutherford Ice Stream Antarctica. *J Glaciol.* 2007;53: 665–672.
13. King EC, Hindmarsh RCA, Stokes CR. Formation of mega-scale glacial lineations observed beneath a west Antarctic ice stream. *Nat Geosci.* 2009;2: 585–596.
14. Smith AM, Murray T. Bedform topography and basal conditions beneath a fast-flowing West Antarctic ice stream. *Quat Sci Rev.* 2009;28: 584–596.
15. Alley RB. In search of ice-stream sticky spots. *J Glaciol.* 1993;39: 447–454.
16. Clark CD, Hughes ALC, Greenwood SL, Spagnolo M, Ng FSL. Size and shape characteristics of drumlins, derived from a large sample, and associated scaling laws. *Quat Sci Rev.* 2009;28:

- 1009 677–692. doi:10.1016/j.quascirev.2008.08.035
- 1010 17. Kleman J, Borgström I. Reconstruction of palaeo-ice sheets: The use of geomorphological data.  
1011 Earth Surf Proc Land. 1996;21: 893–909.
- 1012 18. Duran O, Schwammle V, Lind PG, Herrmann HJ. Size distribution and structure of Barchan  
1013 dune fields. Nonlin Process Geophys. 2011;18: 455–467. doi:10.5194/npg-18-455-2011
- 1014 19. Hillier JK, Smith MJ, Clark CD, Stokes CR, Spagnolo M. Subglacial bedforms reveal an  
1015 exponential size-frequency distribution. Geomorphology. 2013;190: 82–91.  
1016 doi:10.1016/j.geomorph.2013.02.017
- 1017 20. Hindmarsh RCA. Drumlinization and drumlin-forming instabilities: viscous till mechanisms. J  
1018 Glaciol. 1998;44: 293–314.
- 1019 21. Shaw J. Drumlin formation related to inverted melt-water erosional marks. J Glaciol. 1983;29:  
1020 461–479.
- 1021 22. Fowler AC, Spagnolo M, Clark CD, Stokes CR, Hughes ALC, Dunlop P. On the size and shape  
1022 of drumlins. Int J Geomath. 2013;4: 155–165. doi:10.1007/s13137-013-0050-0
- 1023 23. Bak P. How Nature Works: the Science of Self-Organized Criticality. New York: Springer;  
1024 1996.
- 1025 24. Tebbens SF, Burroughs SM, Barton CC, Naar DF. Statistical self-similarity of hotspot seamount  
1026 volumes modeled as self-similar criticality. Geophys Res Lett. 2001;28: 2711–2714.
- 1027 25. Einstein HA. Bedload transport as a probability problem. In: Shen WH, editor. Sedimentation.  
1028 Colorado State University, Fort Collins; 1937. pp. C1–C105.
- 1029 26. van der Mark CF, Blom A, Hulscher SJMH. Quantification of variability in bedform geometry.  
1030 J Geophys Res. 2008;113: F03020. doi:10.1029/2007JF000940
- 1031 27. Singh A, Lanzoni S, Wilcock PR. Multiscale statistical characterization of migrating bed forms  
1032 in gravel and sand rivers. Water Resour Res. 2011;47: W12526. doi:10.1029/2010WR010122
- 1033 28. Fredsøe J. The stability of a sandy river bed. In: Nakato T, Ettema R, editors. Issues and  
1034 Directions in Hydraulics. Balkema, Rotterdam; 1996. pp. 99–114.
- 1035 29. McElroy B, Mohrig D. Nature of deformation of sandy bed forms. J Geophys Res. 2009;144:  
1036 F00A04. doi:10.1029/2008JF001220
- 1037 30. Coleman SE, Nikora VI. Fluvial dunes: initiation, characterization, flow structure. Earth Surf  
1038 Process Landforms. 2011;36: 39–57.
- 1039 31. Spagnolo M, Clark CD, Hughes ALC. Drumlin relief. Geomorphology. 2012;153–154: 179–  
1040 191.
- 1041 32. Hillier JK, Smith M. Testing 3D landform quantification methods with synthetic drumlins in a  
1042 real DEM. Geomorphology. 2012;153: 61–73. doi:doi:10.1016/j.geomorph.2012.02.009
- 1043 33. Hooke R, Medford A. Are drumlins a product of thermo-mechanical instability? Quat Res.  
1044 2013; doi:10.1016/j.yqres.2012.12.002
- 1045 34. Schoof C, Clarke GKC. A model for spiral flows in basal ice and the formation of subglacial  
1046 flutes based on a Reiner-Rivlin rheology for glacial ice. J Geophys Res. 2008;113: B05204.  
1047 doi:10.1029/2007JB004957
- 1048 35. Spagnolo M, Clark CD, Ely JC, Stokes CR, Anderson JB, Andreassen K, et al. Size, shape and  
1049 spatial arrangement of mega-scale glacial lineations from a large and diverse dataset. Earth Surf  
1050 Proc and Landforms. 2014;39: 1432–1448.
- 1051 36. Dowling PF, Spagnolo M, Moller P. Morphometry and core type of streamlined bedforms in

- 1052 southern Sweden from high resolution LiDAR. *Geomorphology*. 2015;236: 54–63.  
 1053 doi:10.1016/j.geomorph.2015.02.018
- 1054 37. Rose J. Glacier sediment patterns and stress transfer associated with the formation of  
 1055 superimposed flutes. *Sediment Geol.* 1989;62: 151–176.
- 1056 38. Goldstein B. Drumlins of the Puget Lowland, Washington state, USA. *Sediment Geol.* 1994;91:  
 1057 299–311.
- 1058 39. Colgan P, Mickelson DM. Genesis of streamlined landforms and flow history of the Green Bay  
 1059 Lobe, Wisconsin, USA. *Sediment Geol.* 1997;111: 7–25.
- 1060 40. Johnson MD, Schomacker A, Benediktsson IO, Geiger AJ, Ferguson A, Ingolfsson O. Active  
 1061 drumlin field revealed at the margin of Mulajokull, Iceland: A surge-type glacier. *Geology*.  
 1062 2010;38: 943–946. doi:10.1130/G31371.1
- 1063 41. Eyles N. Rock drumlins and megaflutes of the Niagara Escarpment, Ontario, Canada: a hard bed  
 1064 landform assemblage cut by the Saginaw-Huron Ice Stream. *Quat Sci Rev.* 2012;55: 34–49.  
 1065 doi:10.1016/j.quascirev.2012.09.001
- 1066 42. Stokes CR, Spagnolo M, Clark CD. The composition and internal structure of drumlins:  
 1067 complexity, commonality, and implications of a unifying theory of their formation. *Earth Sci*  
 1068 *Rev.* 2011;107: 398–422. doi:10.1016/j.earscirev.2011.05.001
- 1069 43. Patterson CJ, Hooke R. Physical environment of drumlin formation. *J Glaciol.* 1995;41: 30–38.
- 1070 44. Clark CD. Emergent drumlins and their clones: from till dilatancy to flow instabilities. *J*  
 1071 *Glaciol.* 2010;51: 1011–1025.
- 1072 45. Stokes CR, Fowler AC, Clark CD, Hindmarsh RCA, Spagnolo M. The instability theory of  
 1073 drumlin formation and its explanation of their varied composition and internal structure. *Quat*  
 1074 *Sci Rev.* 2013;62: 77–96.
- 1075 46. Boyce J, Eyles N. Drumlins carved by deforming till streams below the Laurentide ice sheet.  
 1076 *Geology.* 1991;19: 787–790.
- 1077 47. Hindmarsh RCA. The stability of a viscous till sheet coupled with ice flow, considered at  
 1078 wavelengths less than the ice thickness. *J Glaciol.* 1998;44: 285–292.
- 1079 48. Chapwanya M, Clark CD, Fowler AC. Numerical computations of a theoretical model of ribbed  
 1080 moraine formation. *Earth Surf Proc Land.* 2011;36: 1105–1112.
- 1081 49. Haschenberger J. A probability model of scour and fill depths in gravel-bed channels. *Water*  
 1082 *Resour Res.* 1999;35: 2857–2869.
- 1083 50. Shaw J. The meltwater hypothesis for subglacial bedforms. *Quat Int.* 2002;90: 5–22.
- 1084 51. Ó Cofaigh C, Dowdeswell JA, King EC, Anderson JB, Evans J, Larter RD. Comment on Shaw  
 1085 J, Pugin, A. and Young, R. (2008): A meltwater origin for Antarctic shelf bedforms with special  
 1086 attention to megalineations. *Geomorphology.* 2009;102: 365–375.  
 1087 doi:10.1016/j.geomorph.2009.09.036
- 1088 52. Smalley I, Unwin D. The formation and shape of drumlins and their distribution and orientation  
 1089 in drumlin fields. *J Glaciol.* 1968;7: 377–390.
- 1090 53. Hill AR. The distribution of drumlins in County Down, Ireland. *Ann Assoc Am Geogr.*  
 1091 1973;63: 226–240.
- 1092 54. Van der Meer JJM, Menzies J, Rose J. Subglacial till: the deforming glacier bed. *Quat Sci Rev.*  
 1093 2003;22: 1659–1685.
- 1094 55. Piotrowski J, Larsen NK, Junge FW. Reflections on soft subglacial beds as a mosaic of  
 1095 deforming and stable spots. *Quat Sci Rev.* 2004;23: 993–1000.

- 1096 56. Shoemaker EM. Subglacial water-sheet floods, drumlins and ice-sheet lobes. *J Glaciol.* 1999;45:  
1097 201–213.
- 1098 57. Jerolmack DJ, Mohrig D. A unified model for subaqueous bed form dynamics. *Water Resour*  
1099 *Res.* 2005;41: W12421. doi:10.1029/2005WR004329
- 1100 58. Engelhardt H, Kamb B. Basal sliding of ice stream B, West Antarctica. *J Glaciol.* 1998;44: 223–  
1101 230.
- 1102 59. Hindmarsh RCA. Ice-stream surface texture, sticky spots, waves and breathers: the coupled  
1103 flow of ice, till and water. *J Glaciol.* 1998;44: 589–614.
- 1104 60. Zwally HJ, Abdalati W, Herring T, Larson K, Saba J, Steffen K. Surface melt-induced  
1105 acceleration of Greenland Ice-sheet flow. *Science* (80- ). 2002;297: 218–222.
- 1106 61. Goodwin ID. The nature and origin of a jökulhlaup near Casey Station. *Antarct J Glaciol.*  
1107 1988;34: 95–101.
- 1108 62. Eastwood E, Nield J, Baas A, Kocurek G. Modelling controls on aeolian dune-field pattern  
1109 evolution. *Sedimentology.* 2011;58: 1391–1406.
- 1110 63. Nye JF. A calculation on the sliding of ice over a wavy surface using a Newtonian viscous  
1111 approximation. *Proc R Soc London Ser A.* 1969;311: 445–467.
- 1112 64. Pattyn F, Perichon L, Aschwanden A, Breuer B, De Smedt B, Gagliardini O, et al. Benchmark  
1113 experiments for higher order and full-Stokes ice sheet models (ISMIP-HOM). *Cryosph.* 2008;2:  
1114 95–108.
- 1115 65. Benn DI, Evans DJA. *Glaciers and Glaciation.* 2nd ed. Oxford, UK: Hodder; 2010.
- 1116 66. Meier MF, Post AS. What are glacier surges? *Can J Earth Sci.* 1969;6: 807–819.
- 1117 67. Krimmel RM, Vaughn BN. Columbia Glacier, Alaska: changes in velocity 1977–1986. *J*  
1118 *Geophys Res.* 1987;92: 8961–8.
- 1119 68. Sharp MJ. Surging glaciers: behaviour and mechanisms. *Prog Phys Geogr.* 1988;12: 349–370.
- 1120 69. Willis IC. Inter-annual variations in glacier motion: a review. *Prog Phys Geogr.* 1995;19: 61–  
1121 106.
- 1122 70. Copland L, Sharp MJ, Nienow PW. Links between short-term velocity variations and the  
1123 subglacial hydrology of a predominantly cold polythermal glacier. *J Glaciol.* 2003;49: 337–348.
- 1124 71. Luckman A, Murray T. Seasonal changes in velocity before retreat of Jakobshavn Isbrae,  
1125 Greenland. *Geophys Res Lett.* 2005;32. doi:10.1029/2005GL022519
- 1126 72. MacGregor KR, Riihimäki CA, Anderson RS. Spatial and temporal evolution of rapid basal  
1127 sliding on Bench Glacier, Alaska, USA. *J Glaciol.* 2005;51: 49–63.
- 1128 73. Sigurdsson O, Johnsson T, Johannesson T. Relation between glacier termini and summer  
1129 temperature variations in Iceland since 1930. *Ann Glaciol.* 2007;40: 170–176.
- 1130 74. Dowdeswell JA, Unwin B, Nuttall A, Wingham D. Velocity structure, flow instability and mass  
1131 flux on a large Arctic ice cap from satellite radar interferometry. *Earth Planet Sci Lett.*  
1132 1999;167: 131–140.
- 1133 75. Gravenor C. The origin of drumlins. *Am J Sci.* 1953;251: 674–681.
- 1134 76. Hart JK. The relationship between drumlins and other forms of subglacial glaciotectionic  
1135 deformation. *Quat Sci Rev.* 1997;16: 93–107.
- 1136 77. Schoof C. Cavitation on Deformable Glacier Beds. *SIAM J Appl Math.* 2007;67: 1633–1653.  
1137 doi:10.1137/050646470



- 1138 78. Fowler AC. Instability modelling of drumlin formation incorporating lee-side cavity growth.  
1139 Proc R Soc A Math Phys Eng Sci. 2009;465: 2681–2702. doi:10.1098/rspa.2008.0490
- 1140 79. Dunlop P, Clark CD, Hindmarsh RCA. Bed Ribbing Instability Explanation: Testing a  
1141 numerical model of ribbed moraine formation arising from coupled flow of ice and subglacial  
1142 sediment. J Geophys Res. 2008;113. doi:10.1029/2007JF000954
- 1143 80. Venditti JG, Church MA. Bed form initiation from a flat sand bed. J Geophys Res. 2005;110:  
1144 F01009. doi:10.1029/2004JF000149
- 1145 81. Venditti JG. Bedforms in Sand-Bedded Rivers. In: Shroder J, Wohl E, editors. Treatise on  
1146 Geomorphology. 2013. pp. 137–162.
- 1147 82. Cui Y, Parker G. Sediment pulses in mountain rivers: 2. Comparison between experiments and  
1148 numerical predictions. Water Resour Res. 2003;39: 1240. doi:10.1029/2002WR001805
- 1149 83. Chaytor JD, Ten Brink U, Solow AR, Andrews BD. Size distribution of submarine landslides  
1150 along the US Atlantic margin. Mar Geol. 2009;264: 16–27.
- 1151 84. Leemis LM, McQueston JT. Univariate distribution relationships. Am Stat. 2008;62: 45–53.
- 1152 85. Clark CD. Mega-scale glacial lineations and cross-cutting ice-flow landforms. Earth Surf  
1153 Process Landforms. 1993;18: 1–29.
- 1154 86. Bo T, Zheng X. The formation and evolution of aeolian dune fields under unidirectional wind.  
1155 Geomorphology. 2011;134: 408–416. doi:10.1016/j.geomorph.2011.07.014
- 1156 87. Moll JR, Schilperoort T, De Leeuw JA. Stochastic analysis of bedform dimensions. J Hydraul  
1157 Res. 1987;25: 465–479.
- 1158 88. Øksendal B. Stochastic differential equations: An introduction with applications. 6th ed.  
1159 London: Springer; 2003.
- 1160 89. Mao X. Stochastic differential equations and applications. 2nd ed. Cambridge: Woodhead  
1161 Publishing; 2011.
- 1162 90. Watkins NW. Bunched black (and grouped grey) swans: Dissipative and non-dissipative models  
1163 of correlated extreme fluctuations in complex geosystems. Geophys Res Lett. 2013;40: 1–9.  
1164 doi:10.1002/GRL.50103
- 1165 91. Gregorio A, Iacus SM. Least-squares least-point estimation for the telegraph process observed  
1166 at discrete times. Statistics (Ber). 2010;45: 349–359.
- 1167 92. Vitolo R, Stephenson DS, Cook I, Mitchell-Wallace K. Serial clustering of intense European  
1168 storms. Meteorol Zeitschrift. 2009;18: 411–424. doi:10.1127/0941-2948/2009/0393
- 1169 93. Soong TT. Fundamentals of probability and statistics for engineers. Chichester, UK: Wiley;  
1170 2004.
- 1171 94. Grigoriu M. Stochastic calculus, applications in science and engineering. Boston: Birkhauser;  
1172 2002.
- 1173 95. Rubinstein RY, Kroese DP. Simulation and the Monte Carlo method. 2nd ed. Wiley; 1981.
- 1174 96. Einstein A. Investigations on the theory of brownian motion. 2nd ed. Furth R, editor. Dover,  
1175 London; 1956.
- 1176 97. Tuckwell HC. Elementary Applications of Probability Theory. 2nd ed. London: Chapman and  
1177 Hall; 1995.
- 1178 98. Ross SM. Introduction to Probability Models. 5th ed. Academic Press; 1993.
- 1179 99. Kougioumtzoglou IA, Spanos PD. An analytical Wiener path integral technique for non-  
1180 stationary response determination of nonlinear oscillators. Probabilistic Eng Mech. 2012;28:

125–131.

100. Ross SM. Stochastic Processes. 2nd ed. Wiley; 1996.

101. Grimmett G, Stirzaker D. Probability and Random Processes. Oxford, UK: OUP; 2001.

102. Hillier JK, Smith MJ, Armugam R, Barr, Boston C, Clark CD, et al. Manual mapping of drumlins in synthetic landscapes to assess operator effectiveness. *J Maps*. 2014;11: 719–729. doi:doi:10.1080/17445647.2014.957251103. Smith MJ, Rose J, Booth S. Geomorphological mapping of glacial landforms from remotely sensed data: an evaluation of the principal data sources and an assessment of their quality. *Geomorphology*. 2006;76: 148–165.104. Hollingsworth SE. The glaciation of western Edenside and adjoining areas and the drumlins of Edenside and the Solway basin. *Quart J Geol Soc London*. 1931;87: 281–359.

105. Wilks DS. Statistical methods in the atmospheric sciences. 2ns ed. London: Academic Press; 2006.

106. Gorini C. Master Math: Probability. 1st ed. Boston, USA: Cengage Learning; 2012.

107. Schoof C. Mathematical models of glacier sliding and drumlin formation [Internet]. Oxford University. 2002. Available: <http://www.maths.ox.ac.uk/research/theses>108. Rose J. Drumlins as part of a glacier bedform continuum. In: Menzies J, Rose J, editors. *Drumlin Symposium*. Balkema, Rotterdam; 1987. pp. 103–116.109. Aario R. Classification and terminology of moranic landforms in Finland. *Boreas*. 1977;6: 77–100.110. Fowler AC. An instability mechanism for drumlin formation. In: Maltman AJ, Hubbard B, Hambrey MJ, editors. *Deformation of Glacial Materials*. Geological. London: Geol. Soc. Publishing House; 2000. pp. 307–319.111. Hubbard AL, Bradwell T, Golledge N, Hall A, Patton H, Sugden D, et al. Dynamic cycles, ice streams and their impact on the extent, chronology and deglaciation of the British-Irish ice sheet. *Quat Sci Rev*. 2009;28: 758–776.

112. Turcotte DL. Fractals and Chaos in Geology and Geophysics. 2nd ed. Cambridge: Cambridge University Press; 1997.

113. Stokes CR, Spagnolo M, Clark CD, Ó Cofaigh C, Lian OB, Dunstone RB. Formation of mega-scale glacial lineations on the Dubawnt Lake Ice Stream bed: 1. Size, shape and spacing from a large remote sensing dataset. *Quat Sci Rev*. 2013;

114. Mandlebrot BB. The Fractal Geometry of Nature. W. H. Freeman and Company, New York; 1983.

115. Mendenhall W, Wackerly DD, Scheaffer RL. Mathematical statistics with applications. 4th ed. Belmont, California: Duxbury Press; 1990.

116. Shen H, Brown LD, Zhi H. Efficient estimation of log-normal means with application to pharmacokinetic data. *Stat Med*. 2006;25: 3023–3038. doi:10.1002/sim.2456

## Supporting information

**S1 File. Zip file, containing data and worked examples of parameter calculation in EXCEL sheets, and a README file explaining its contents.**

1222

1223